



Simon, N., Kusakina, J., Fernandez-Lopez, A., Chembath, A., Belbin, F., & Dodd, A. (2018). The Energy-Signaling Hub SnRK1 Is Important for Sucrose-Induced Hypocotyl Elongation. *Plant Physiology*, 176, 1299-1310. <https://doi.org/10.1104/pp.17.01395>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1104/pp.17.01395](https://doi.org/10.1104/pp.17.01395)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via ASPB at <http://www.plantphysiol.org/content/176/2/1299> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>

The energy-signalling hub SnRK1 is important for sucrose-induced hypocotyl elongation

Noriane M. L. Simon^(1, 5), Jelena Kusakina^(2, 5), Ángela Fernández-López⁽¹⁾, Anupama Chembath⁽³⁾,
Fiona E. Belbin⁽¹⁾, Antony N. Dodd^{(1, 4)*}

1. School of Biological Sciences, University of Bristol, Life Sciences Building, 24 Tyndall
Avenue, Bristol BS8 1TQ, U.K.

2. Present address: Centre for Plant Sciences, Faculty of Biological Sciences, University of Leeds,
Leeds LS2 9JT, U.K.

3. Present address: Institute for Cell and Molecular Biosciences, Faculty of Medical Sciences,
Newcastle University, Framlington Place, Newcastle upon Tyne NE2 4HH, U.K.

4. Cabot Institute, University of Bristol, Bristol BS8 1UJ, U.K.

5. Equal contribution.

***Corresponding author:** Antony Dodd, School of Biological Sciences, University of Bristol, Life
Sciences Building, 24 Tyndall Avenue, Bristol BS8 1TQ, U.K
antony.dodd@bristol.ac.uk, tel +44 (0)117 394 1176.

Word counts: Summary: 211 Introduction: 1035 Methods: 600 Results: 2366 Discussion: 1668
Acknowledgements 113 (Main text total: 5669). Figure legends: 666.

Author Contributions: A.D. and J.K. conceived the study; N.S, J.K., A.F.L., F.B. and A.C.
performed experiments; N.S., J.K., A.F.L. and A.D. analyzed data; N.S., J.K. and A.D. interpreted
data and wrote the paper.

One-sentence summary: Sucrose-induced hypocotyl elongation involves the energy-signalling
hub SnRK1.

Running title: Sucrose-induced hypocotyl elongation

30 **Abstract**

31 Emerging seedlings respond to environmental conditions such as light and temperature to optimize
32 their establishment. Seedlings grow initially through elongation of the hypocotyl, which is
33 regulated by signalling pathways that integrate environmental information to regulate seedling
34 development. The hypocotyls of *Arabidopsis thaliana* also elongate in response to sucrose. Here,
35 we investigated the role of cellular sugar-sensing mechanisms in the elongation of hypocotyls in
36 response to sucrose. We focused upon the role of SnRK1, which is a sugar-signalling hub that
37 regulates metabolism and transcription in response to cellular energy status. We also investigated
38 the role of TPS1, which synthesizes the signalling sugar trehalose-6-phosphate (Tre6P) that is
39 proposed to regulate SnRK1 activity. Under light/dark cycles, we found that sucrose-induced
40 hypocotyl elongation did not occur in *tps1* mutants and overexpressors of KIN10
41 (AKIN10/SnRK1.1), a catalytic subunit of SnRK1. We demonstrate that the magnitude of sucrose-
42 induced hypocotyl elongation depends on the day length and light intensity. We identified roles for
43 auxin and gibberellin signalling in sucrose-induced hypocotyl elongation under short photoperiods.
44 We found that sucrose-induced hypocotyl elongation under light/dark cycles does not involve
45 another proposed sugar sensor, HEXOKINASE1, or the circadian oscillator. Our study identifies
46 novel roles for KIN10 and TPS1 in mediating a signal that underlies sucrose-induced hypocotyl
47 elongation in light/dark cycles.

48

49

50 **Introduction**

51 Emerging seedlings monitor the environment to optimize their establishment and out-compete
52 neighbouring plants (Salter et al., 2003; Weinig et al., 2007; Koini et al., 2009; Keuskamp et al.,
53 2010; Crawford et al., 2012). Seedlings grow initially through cell expansion within the hypocotyl,
54 which elongates rapidly to optimize light capture by the cotyledons. Hypocotyl elongation is
55 controlled by several signalling pathways that converge upon phytohormones to regulate cell
56 expansion (Lincoln et al., 1990; Collett et al., 2000). Examples of signals that adjust hypocotyl
57 elongation include phytochrome-mediated signals concerning the ratio of red to far red light
58 (R:FR) (Casal, 2013), blue light (Liscum and Hangarter, 1991), UV-B light (Kim et al., 1998;
59 Hayes et al., 2014), temperature (Koini et al., 2009; Wigge, 2013; Mizuno et al., 2014),
60 photoperiod and the circadian oscillator (Dowson-Day and Millar, 1999; Más et al., 2003; Nusinow
61 et al., 2011). These signals are integrated by the PHYTOCHROME INTERACTING FACTOR
62 (PIF)-family of basic helix-loop-helix transcription factors. The PIFs are signalling hubs that
63 control plant development through genome-wide transcriptional alterations. One outcome of these
64 PIF-mediated transcriptional changes are the alterations in phytohormone signalling that regulate
65 hypocotyl elongation (Lorrain et al., 2008; Leivar and Quail, 2011).

66 Hypocotyl length is also increased by exogenous and endogenous sugars (Kurata and Yamamoto,
67 1998; Takahashi et al., 2003; Zhang et al., 2010; Liu et al., 2011; Stewart et al., 2011; Stewart
68 Lilley et al., 2012; Zhang et al., 2015; Zhang et al., 2016). Under light/dark cycles, exogenous
69 sugars are proposed to cause hypocotyl elongation by inducing auxin signals through the PIF-
70 mediated gene regulation (Stewart et al., 2011; Stewart Lilley et al., 2012). Under extended
71 darkness, brassinosteroid and GA phytohormones are involved in sugar-induced hypocotyl
72 elongation, which may also involve the target of rapamycin (TOR) kinase regulator of energy- and
73 nutrient-responses (Zhang et al., 2010; Dobrenel et al., 2011; Zhang et al., 2015; Zhang et al.,

2016). This elongation phenotype in darkness is thought to form a response to the starvation
 conditions that arise when plants are cultivated under periods of darkness exceeding the length of
 the daily light/dark cycle (Graf et al., 2010; Zhang et al., 2016). In comparison to these known
 roles for phytohormones and transcriptional regulators, the contribution of sugar sensing
 mechanisms to sucrose-induced hypocotyl elongation remain unknown.
 Several sugar- or energy-signalling mechanisms underlie the metabolic and developmental
 responses of plants to sugars. One mechanism involves the Sucrose non-fermenting 1 (Snf1)-
 related protein kinase SnRK1 (Baena-González et al., 2007; Baena-González and Sheen, 2008),
 and another involves HEXOKINASE1 (Jang et al., 1997; Moore et al., 2003). SnRK1 controls
 metabolic enzymes directly by protein phosphorylation (Baena-González and Sheen, 2008). It also
 regulates > 1000 transcripts in response to carbohydrate availability, for example by adjusting bZIP
 transcription factor activity (Baena-González et al., 2007; Smeekens et al., 2010; Delatte et al.,
 2011; Mantioli et al., 2011; Mair et al., 2015). Both SnRK1- and hexokinase-mediated sugar
 signalling involve specific sugars functioning as signalling molecules that provide cellular
 information concerning sugar availability. For example, SnRK1 activity is thought to be regulated
 by trehalose-6-phosphate (Tre6P), whose concentration tracks the cellular concentration of sucrose
 (Lunn et al., 2006; Zhang et al., 2009; Nunes et al., 2013; Yadav et al., 2014). Tre6P is synthesized
 from UDP glucose and glucose-6-phosphate, which are derived from mobilized and transported
 sucrose, and also directly from photosynthesis. In *Arabidopsis thaliana*, Tre6P is
 synthesized by trehalose-6-phosphate synthase (TPS). Of 11 TPS homologs encoded by the
Arabidopsis genome, TREHALOSE-6-PHOSPHATE SYNTHASE1 (TPS1) synthesizes Tre6P in
 plants (Gómez et al., 2010; Vandesteene et al., 2010), and TPS2 and TPS4 are catalytically active
 in yeast complementation assays (Delorge et al., 2015). Tre6P is believed to regulate SnRK1-
 mediated signalling by suppressing the activity of SNF1-RELATED PROTEIN KINASE1.1

(KIN10/AKIN10/SnRK1.1), which is a catalytic subunit of SnRK1 that is fundamental to the signalling role of SnRK1 (Baena-González et al., 2007; Zhang et al., 2009; Nunes et al., 2013). Manipulation of Tre6P metabolism in plants alters developmental phenotypes. For example, *tps1* knockout mutants undergo seedling developmental arrest (Gómez et al., 2006), expression of bacterial Tre6P synthase (*otsA*) or phosphatase (*otsB*) affects leaf senescence (Wingler et al., 2012), and Tre6P and KIN10 act within a photoperiod-response pathway that controls the induction of flowering (Baena-González et al., 2007; Gómez et al., 2010; Wahl et al., 2013). Signalling by Tre6P and KIN10 is also important for the regulation of growth rates. Growth is increased by sucrose in the presence of Tre6P (Schluepmann et al., 2003; Paul et al., 2010), but the lack of a quantitative (correlative) relationship between relative growth rates and [Tre6P] suggests that a threshold [Tre6P] is required for growth to occur (Nunes et al., 2013). Therefore, it has been suggested that control of KIN10/11 by [Tre6P] may ‘prime’ the regulation of growth-related genes to capitalize upon increased energy availability, rather than by inducing growth directly (Nunes et al., 2013). Remarkably, the impact of this pathway is sufficiently global that its manipulation can increase maize yields by almost 50% (Nuccio et al., 2015) and increase the yield and drought tolerance of wheat (Griffiths et al., 2016). Given the importance of Tre6P metabolism and SnRK1 for growth regulation under cycles of light and dark, we wished to determine whether this energy-signalling mechanism is important for the regulation of sucrose-induced hypocotyl elongation. Moreover, because Tre6P signalling is reported to act upon GA and auxin signalling genes (Paul et al., 2010; Li et al., 2014) and these phytohormones are involved in sucrose-induced hypocotyl elongation (Zhang et al., 2010; Stewart Lilley et al., 2012), we reasoned that SnRK1 might act upon these phytohormones to regulate sucrose-induced hypocotyl elongation.

121 Here, we identified a novel role for Tre6P and KIN10 in the mechanisms that cause sucrose-
122 induced hypocotyl elongation. We focused upon light/dark cycles rather than conditions of
123 extended darkness (Zhang et al., 2010; Zhang et al., 2015; Zhang et al., 2016), because we wished
124 to identify mechanisms that regulate growth and development under regimes more representative
125 of real-world growing conditions that do not elicit prolonged starvation. We found that the
126 sensitivity of hypocotyl elongation to sugars depends on the photoperiod and light intensity. We
127 identified that KIN10 is important for expression of transcripts encoding auxin-induced expansins.
128 Our data reveal a new mechanistic link between carbohydrate supply, energy sensing and
129 phytohormone signalling during seedling emergence.

130 **Results**

131 *KIN10 and TPS1 are required for sucrose-induced hypocotyl elongation in light/dark cycles*

132 We investigated whether KIN10 and TPS1 contribute to sucrose-induced hypocotyl elongation
133 under light/dark cycles (Kurata and Yamamoto, 1998; Takahashi et al., 2003; Stewart et al., 2011;
134 Stewart Lilley et al., 2012). We studied hypocotyl elongation in transgenic Arabidopsis where
135 KIN10 activity was manipulated by overexpressing the catalytic subunit of KIN10 (KIN10-ox)
136 (Baena-González et al., 2007). Although KIN10 activity is regulated post-translationally by Tre6P
137 (Zhang et al., 2009), KIN10 overexpression alone alters the abundance of energy-response
138 transcripts in protoplasts (Baena-González et al., 2007). We used KIN10 overexpression rather
139 than knockouts, because KIN10/11 double knockouts disrupt pollen production and are lethal
140 (Zhang et al., 2001; Baena-González et al., 2007). We also used hypomorphic TILLING (targeted
141 induced local lesions in genomes) mutants with reduced TPS1 activity (*tps1-11*, *tps1-12*) (Gómez
142 et al., 2006; Gómez et al., 2010), which is preferable to *tps1* loss-of-function mutants that cause
143 seedling developmental arrest (Gómez et al., 2006).

144 First, we investigated the effect of exogenous sucrose upon hypocotyl elongation in a variety of
145 photoperiods (Fig. 1). Under 4 h and 8 h photoperiods, sucrose supplementation of wild type
146 seedlings caused a significant increase in hypocotyl length relative to the sorbitol control (2.1-fold
147 and 2.3-fold relative to sorbitol controls, under 4 h and 8 h photoperiods respectively) (Fig. 1A-E).
148 In comparison, under 16 h photoperiods and constant light conditions exogenous sucrose did not
149 promote hypocotyl elongation (Fig. 1A-E).

150 Next, we investigated roles of KIN10 in sucrose-induced hypocotyl elongation under light/dark
151 cycles. Under 8 h photoperiods, the hypocotyls of two KIN10-ox lines (Baena-González et al.,
152 2007) did not elongate significantly in response to exogenous sucrose relative to the MS control
153 (Fig. 1B). Both KIN10-ox lines elongated 1.5-fold in response to sucrose relative to the sorbitol
154 control (Fig. 1B). Exogenous sucrose caused no significant increase in the hypocotyl length of
155 KIN10-ox seedlings under 4 h photoperiods (Fig. 1C). Hypocotyls of the *L. er.* background and
156 KIN10-ox appeared shorter when supplemented with exogenous sucrose in constant light and 16 h
157 photoperiods. However, this could be an osmotic effect rather than a sucrose response because
158 hypocotyl elongation responded identically to sucrose and the sorbitol control (Fig. 1B).

159 Since KIN10 activity is thought to be regulated by Tre6P (Zhang et al., 2009), we investigated the
160 role of the Tre6P biosynthetic enzyme TPS1 in sucrose-induced hypocotyl elongation under
161 light/dark cycles. In two *tps1* TILLING mutants under 8 h photoperiods, sucrose supplementation
162 caused a significant 2.3-fold increase in hypocotyl length in the wild type relative to the sorbitol
163 control, compared with 1.6-fold and 1.3-fold increases in hypocotyl length in *tps1-11* and *tps1-12*
164 respectively (Fig. 1D). Under 4 h photoperiods, sucrose caused a significant 2-fold increase in
165 hypocotyl length of the wild type relative to the sorbitol control, compared with no significant
166 increase in length in *tps1-11* and a significant 1.5-fold increase in hypocotyl length in *tps1-12* (Fig.
167 1E). Together, these experiments with KIN10 overexpressors and *tps1* mutants indicate that TPS1

168 and KIN10 are involved in one or more mechanisms that increase hypocotyl length in response to
169 exogenous sucrose. This suggests that SnRK1-mediated energy signalling regulates hypocotyl
170 elongation in response to sucrose supplementation.

171 *HEXOKINASE1 is not required for sucrose-induced hypocotyl elongation under light/dark cycles*
172 Hexokinase is thought to function as a sugar sensor that regulates development in response to the
173 concentration of glucose (Jang et al., 1997; Moore et al., 2003), so we investigated whether
174 hexokinase-based signalling also contributes to sucrose-induced hypocotyl elongation. For this, we
175 measured the elongation of hypocotyls in response to exogenous sucrose in the *glucose insensitive2*
176 (*gin2-1*) mutant of HEXOKINASE1. Overall, *gin2-1* hypocotyls were slightly shorter than the wild
177 type under all conditions tested (Fig. 1F). Exogenous sucrose caused a significant increase in
178 hypocotyl length of wild type and *gin2-1* seedlings, producing hypocotyls 63% and 67% longer
179 than the osmotic control in the wild type and *gin2-1*, respectively (Fig. 1F). Therefore, sucrose
180 caused a similar magnitude of hypocotyl elongation in *gin2-1* and the wild type. This suggests that
181 interconversion of sucrose to glucose, and therefore hexokinase-based glucose signalling, does not
182 contribute to sucrose-induced hypocotyl elongation in short photoperiods.

183 *Relationship between day-length, light intensity and sucrose-induced hypocotyl elongation*

184 Our data suggest that the magnitude of the sucrose-induced increase in hypocotyl length depends
185 upon the photoperiod or the quantity of light received. In the wild type, sucrose increased
186 hypocotyl length under short (4 h or 8 h) but not long (16 h or constant light) photoperiods under
187 photosynthetically active radiation (PAR) of 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 1B-E, Fig. 2A). In addition,
188 sucrose caused significantly greater hypocotyl elongation under 4 h photoperiods compared with 8
189 h photoperiods of 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 2A). We reasoned that these varying responses to sucrose
190 might arise from differences in total daily PAR received under each of these conditions, or

alternatively from the sensing of photoperiod length. To investigate this we compared the magnitude of sucrose-induced hypocotyl elongation under the same total daily integrated PAR, under longer photoperiods (16 h at $40 \mu\text{mol m}^{-2} \text{s}^{-1}$ and 8 h at $80 \mu\text{mol m}^{-2} \text{s}^{-1}$) and under shorter photoperiods (8 h at $40 \mu\text{mol m}^{-2} \text{s}^{-1}$ and 4 h at $80 \mu\text{mol m}^{-2} \text{s}^{-1}$). Under a 16 h photoperiod at $40 \mu\text{mol m}^{-2} \text{s}^{-1}$, sucrose caused a significant increase in hypocotyl length (Fig. 2B, C). This contrasts a 16 h photoperiod at $100 \mu\text{mol m}^{-2} \text{s}^{-1}$, where sucrose did not promote hypocotyl elongation (Fig. 1, Fig. 2A). This suggests that the quantity of light received influences the sensitivity of hypocotyl elongation to sucrose. Under 8 h photoperiods, sucrose caused greater hypocotyl elongation under $40 \mu\text{mol m}^{-2} \text{s}^{-1}$ (mean 4.1 mm increase) than under $80 \mu\text{mol m}^{-2} \text{s}^{-1}$ (mean 3.3 mm increase), which also suggests that hypocotyl elongation is more responsive to sucrose under lower light conditions (Fig. 2B, D). When daily integrated PAR was the same under 4 h and 8 h photoperiods, there was no difference in the increase in hypocotyl length caused by sucrose (Fig. 2D, E). These responses suggest that daily integrated PAR influences the magnitude of sucrose-induced hypocotyl elongation. However, the magnitude of sucrose-induced hypocotyl elongation was significantly less under 16 h photoperiods at $40 \mu\text{mol m}^{-2} \text{s}^{-1}$ than 8 h photoperiods at $80 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 2B, C), suggesting that under long photoperiods, the magnitude of sucrose-induced hypocotyl elongation could be also determined by a photoperiod-response mechanism acting independently from daily integrated PAR. These data provide the insight that the photoperiod-sensitivity of sucrose-induced hypocotyl elongation is determined by both the absolute photoperiod and the amount of light received.

Interaction between hypocotyl elongation by exogenous sucrose and the circadian oscillator

The circadian oscillator regulates hypocotyl elongation because the accumulation of PIF proteins is restricted to the end of the night (Nozue et al., 2007; Nusinow et al., 2011). Since the circadian

oscillator responds to exogenous and endogenous sugars (Dalchau et al., 2011; Haydon et al., 2013) and KIN10 overexpression can lengthen circadian period (Shin et al., 2017), we investigated whether sucrose-induced increases in hypocotyl length under short photoperiods involve the circadian oscillator. First, we tested whether the circadian oscillator components CIRCADIAN CLOCK ASSOCIATED1 (CCA1), LATE ELONGATED HYPOCOTYL (LHY) and TIMING OF CAB2 EXPRESSION1 (TOC1) are required for sucrose-induced hypocotyl elongation using the *cca1-11 lhy-21 toc1-21* triple mutant (Ding et al., 2007). *cca1-11 lhy-21 toc1-21* causes circadian arrhythmia under constant light and temperature, and disrupts rhythms of oscillator transcripts, including evening complex components that regulate hypocotyl elongation (Ding et al., 2007). Under 4 h photoperiods, the magnitude of the sucrose-induced increase in hypocotyl length was unaltered in *cca1-11 lhy-21 toc1-21* (Fig. 3A; Fig. S1). Under 4 h photoperiods the hypocotyls of *cca1-11 lhy-21 toc1-21* were of similar length to the wild type (Fig. 3A), whereas under 8 h photoperiods, *cca1-11 lhy-21 toc1-21* has longer hypocotyls than the wild type (Ding et al., 2007). We also investigated whether two proteins that confer sugar sensitivity to the circadian oscillator, GIGANTEA (GI) and PSEUDO-RESPONSE REGULATOR7 (PRR7) (Dalchau et al., 2011; Haydon et al., 2013), contribute to sucrose-induced hypocotyl elongation under short photoperiods. We tested this because the *prp7-11* mutation renders the oscillator insensitive to sugar signals that entrain the oscillator (Haydon et al., 2013), and the *gi-11* mutation alters oscillator responses to long-term exposure to exogenous sucrose (Dalchau et al., 2011). In all cases, *gi-11* had longer hypocotyls than the wild type (Fig. 3B), but the magnitude of the sucrose-induced increase in hypocotyl length was unaltered in *gi-11* relative to the wild type (Fig. 3D). Likewise, the *prp7-11* mutant also did not alter the magnitude of sucrose-induced increases in hypocotyl length (Fig. 3C, D).

237 These experiments indicate that two mechanisms providing sugar inputs to the circadian oscillator
 238 (Dalchau et al., 2011; Haydon et al., 2013) and three core oscillator components do not contribute
 239 to sucrose-induced increases in hypocotyl length under short photoperiods.

240 *Phytohormone signalling and sucrose-induced hypocotyl elongation under light/dark cycles: auxin*
 241 Sucrose-induced hypocotyl elongation in the light involves auxin and GA signalling (Zhang et al.,
 242 2010; Stewart Lilley et al., 2012). We investigated the involvement of phytohormones in sucrose-
 243 induced hypocotyl elongation under light/dark cycles, and their relationship with SnRK1-mediated
 244 signalling. First, we examined the effect of the inhibitor of polar auxin transport 1-N-
 245 naphthylphthalamic acid (NPA) upon sucrose-induced hypocotyl elongation. NPA inhibited sucrose-
 246 induced hypocotyl elongation in a concentration-dependent manner, such that 10 μ M NPA
 247 completely abolished sucrose-induced elongation (Fig. 4A). Consistent with previous work
 248 (Stewart Lilley et al., 2012), this indicates that under light/dark cycles sucrose-induced hypocotyl
 249 elongation is auxin-dependent. Next, we examined the responses of auxin- and PIF-dependent
 250 expansin transcripts to sucrose. Expansins are a large family of cell-wall modifying enzymes that
 251 allow turgor-driven cell expansion, and some expansin transcripts are upregulated by auxins in a
 252 PIF-dependent manner during hypocotyl elongation (Li et al., 2002; Miyazaki et al., 2016;
 253 Gangappa and Kumar, 2017). We examined *EXPANSIN A4 (EXPA4)*, *EXPA8* and *EXPA11*
 254 transcripts, which are auxin-induced in seedlings (Goda et al., 2004; Esmon et al., 2006; Winter et
 255 al., 2007; Lee et al., 2009). *EXPA8* and *EXPA11* transcripts were upregulated by conditions of
 256 constant darkness, which also increases hypocotyl elongation (Fig. S2A) (Boylan and Quail, 1991),
 257 and downregulated by 10 μ M NPA, which suppresses hypocotyl elongation (Fig. S2B) (Stewart
 258 Lilley et al., 2012). *EXPA4* was unaltered by these conditions (Fig. S2). Therefore, *EXPA8* and
 259 *EXPA11* transcript abundance was increased by conditions that promote hypocotyl elongation, and

260 reduced by conditions that suppress hypocotyl elongation. Next, we monitored the change in
261 abundance of these two expansin transcripts in response to sucrose under 4 h photoperiods. In the
262 wild type, *EXPA11* transcripts were upregulated by 3% sucrose, whereas *EXPA8* transcripts were
263 not upregulated by sucrose relative to the controls (Fig. 4B-E). In KIN10-ox, where sucrose does
264 not promote hypocotyl elongation under light/dark cycles, *EXPA8* and *EXPA11* transcripts were
265 not increased by sucrose (Fig. 4B-E). *EXPA8* was sucrose-induced relative to the controls in *tps1*-
266 11, but not in *tps1*-12 (Fig. 4B, C). *EXPA11* transcripts were sucrose-induced in both *tps1*-11 and
267 *tps1*-12 (Fig. 4D, E). The induction of these two expansin transcripts by sucrose in *tps1* mutants
268 was unexpected, because both KIN10-ox and *tps* mutants suppress sucrose-induced hypocotyl
269 elongation under short photoperiods (Fig. 1). We also examined several other transcripts associated
270 with auxin biosynthesis or responses, but the osmotic controls caused substantial alterations in
271 transcript abundance that prevented interpretation of their regulation by sucrose (Fig. S3).

272 *Phytohormone signalling and sucrose-induced hypocotyl elongation under light/dark cycles:*
273 *gibberellins*

274 We tested whether GA signalling also contributes to sucrose-induced hypocotyl elongation under
275 short photoperiods. After germination, wild type seedlings were transferred to media containing
276 3% sucrose or an osmotic control, supplemented with combinations of the GA biosynthesis
277 inhibitor paclobutrazol (PAC), GA, or a carrier control. Consistent with previous studies, wild type
278 seedlings grown on media supplemented with PAC or PAC and GA had significantly shorter
279 hypocotyls than controls (Fig. 5A) (Cowling and Harberd, 1999; Liu et al., 2011). PAC abolished
280 sucrose-induced hypocotyl elongation, with a small hypocotyl length rescue occurring when GA
281 was supplied in combination with PAC (Fig. 5A). We confirmed that the GA was active by

282 demonstrating that, consistent with previous reports (Cowling and Harberd, 1999), hypocotyl
283 length is increased by GA supplementation (Fig. S4).

284 GA increases growth by causing degradation of DELLA growth repressor proteins, and also
285 through DELLA-independent mechanisms (Peng et al., 1997; Fu et al., 2002; Cheng et al., 2004;
286 Cao et al., 2006). Therefore, we investigated the involvement of DELLA proteins in sucrose-
287 induced hypocotyl elongation under light/dark cycles. The *gai-1* mutant harbours a deletion within
288 the DELLA domain of *GIBBERELLIC ACID INSENSITIVE* (*GAI*), which prevents GA-induced
289 proteasomal degradation of *GAI* (Peng et al., 1997; Fu et al., 2002). Under 4 h photoperiods,
290 sucrose supplementation increased hypocotyl length in *gai-1*, but the magnitude of sucrose-induced
291 elongation in *gai-1* was reduced compared with the wild type (hypocotyls became 36.5% longer in
292 *gai-1* in response to sucrose, compared with 59.2% longer in the wild type) (Fig. 5B). Under 16 h
293 photoperiods, sucrose did not induce hypocotyl elongation in the wild type or *gai-1* (Fig. 5B),
294 which is consistent with Fig. 1B, C. We also examined the effect of a mutant lacking all five
295 DELLA proteins upon sucrose-induced hypocotyl elongation under light/dark cycles (Koini et al.,
296 2009). Under short photoperiods, sucrose-induced hypocotyl elongation was unaltered in this
297 mutant (Fig. 5C). Interestingly, under long photoperiods sucrose promoted hypocotyl elongation in
298 the DELLA global mutant, whereas sucrose was without effect upon wild type hypocotyls (Fig.
299 5C). The partial attenuation of sucrose-induced hypocotyl elongation in *gai-1* (Fig. 5B) combined
300 with the derepression of sucrose-induced hypocotyl elongation under long photoperiods in the
301 DELLA global mutant (Fig. 5C) suggests that DELLA-mediated GA signalling contributes to, but
302 does not exclusively control, sucrose-induced hypocotyl elongation.

303 *Phytohormone signalling and sucrose-induced hypocotyl elongation under light/dark cycles:*
304 *abscisic acid*

305 ABA suppresses seedling development (Belin et al., 2009) and several studies have linked Tre6P
306 and abscisic acid (ABA) signalling (Avonce et al., 2004; Ramon et al., 2007; Gómez et al., 2010;
307 Debast et al., 2011). Therefore, we investigated whether ABA signalling contributes to sucrose-
308 induced hypocotyl elongation under light/dark cycles. Sucrose-induced hypocotyl elongation was
309 unaffected by the ABA receptor quadruple mutant *pyr1-1 pyl1-1 pyl2-1 pyl4-1*, which is highly
310 ABA-insensitive (Park et al., 2009) (Fig. S5). This suggests that PYR/PYL-mediated ABA
311 signalling does not participate in the mechanisms underlying sucrose-induced hypocotyl elongation
312 under light/dark cycles.

313 **Discussion**

314 *KIN10 and TPS1 contribute to sugar-induced hypocotyl elongation under light/dark cycles*

315 Here, we make the new finding that a mechanism involving KIN10 activity and Tre6P metabolism
316 regulates sucrose-induced hypocotyl elongation under light/dark cycles. Whilst hypocotyl
317 elongation arises from cell expansion rather than growth through increases in cell number
318 (Gendreau et al., 1997), our data are consistent with studies demonstrating that Tre6P metabolism
319 is a crucial regulator of growth responses to sucrose. For example, Arabidopsis seedlings
320 overexpressing the bacterial Tre6P phosphatase *otsB*, which reduces [Tre6P], accumulate less
321 biomass compared with the wild type when supplemented with sucrose (Schluepmann et al., 2003).
322 The converse is also true; *otsA* (TPS) overexpressors, in which [Tre6P] is increased, accumulate
323 more biomass than the wild type when supplemented with sucrose (Schluepmann et al., 2003).
324 Therefore, our data using *tps1* mutants as a proxy for altered Tre6P metabolism provide new

evidence to support the notion that Tre6P promotes growth under conditions of increased sucrose availability (Schluepmann et al., 2003; Zhang et al., 2009). Overexpression in Arabidopsis of the bacterial Tre6P synthase *otsA* has been reported to produce seedlings having shorter hypocotyls than the wild type (Paul et al., 2010). The sucrose-insensitivity of hypocotyl elongation in *tps1* mutants (Fig. 1) and the shorter hypocotyls in seedlings with increased [Tre6P] (*otsA-ox*) may appear to conflict with each other (Paul et al., 2010). However, the experiments are not directly comparable. We found that exogenous sucrose only caused hypocotyl elongation under short photoperiods or lower light conditions (Fig. 2). In comparison, the *otsA-ox* experiments involved 16 h photoperiods at higher PAR ($150 \mu\text{mol m}^{-2} \text{s}^{-1}$) and shaking liquid culture (Zhang et al., 2009), both of which could mask the hypocotyl elongation response that we investigated. Our experiments suggest that increased KIN10 activity might attenuate the elongation response of hypocotyls to exogenous sucrose under light/dark cycles. The KIN10-ox lines that we used overexpress the catalytic subunit of SnRK1 (Baena-González et al., 2007). KIN10 overexpression downregulates transcripts associated with anabolic processes and upregulates transcripts associated with energy starvation (Baena-González et al., 2007). Therefore, in our experiments KIN10 overexpression may have stopped seedlings from taking advantage of the greater energy availability caused by sucrose supplementation, so preventing sucrose-induced hypocotyl elongation in KIN10-ox (Fig. 1).

Photoperiod-dependency of sugar-induced hypocotyl elongation

We made the new finding that under relatively high light, exogenous sucrose increases hypocotyl length in photoperiods of 8 h and shorter, but not under long photoperiods or constant light (Fig. 1, Fig. 2). These data reconcile differences between previous studies of sucrose-induced hypocotyl

348 elongation. Previous studies reporting sucrose-insensitivity of hypocotyl elongation in the light
349 were conducted in continuous light (Zhang et al., 2010), in which we also found sucrose to be
350 without effect upon hypocotyls (Fig. 1B, Fig. 2A). In comparison, studies reporting that sucrose
351 does promotes hypocotyl elongation in the light were conducted under 8 h photoperiods (Stewart et
352 al., 2011; Stewart Lilley et al., 2012), where we likewise found that sucrose causes hypocotyl
353 elongation (Fig. 1B, Fig. 2). Therefore, the sensitivity of hypocotyls to sucrose-induced elongation
354 depends upon the photoperiod or the amount of light received each day.

355 One explanation for this response could be that the daily quantity of light determines the magnitude
356 of sucrose-induced hypocotyl elongation through the accumulation of photosynthetic metabolites.
357 Our experiments indicate that under shorter photoperiods, the sensitivity of hypocotyl elongation to
358 sucrose depends upon the total amount of daily light (Fig. 2A, D, E). Furthermore, sucrose-induced
359 hypocotyl elongation under long photoperiods only occurred when the seedlings were under lower
360 light conditions (Fig. 2A, B, C). One interpretation is that under long photoperiods and higher light,
361 cells are replete with sugars (Sulpice et al., 2014) therefore supplementation with exogenous
362 sucrose has a relatively small effect upon the hypocotyl length of already sugar-rich seedlings. In
363 contrast, under short photoperiods or lower light the background level of endogenous sugar is
364 lower (Sulpice et al., 2014), so supplementation with exogenous sucrose has a greater effect upon
365 hypocotyl length.

366 An alternative interpretation is that PIFs integrate light signals derived from photoreceptors with
367 SnRK1-mediated sugar signals to modulate the sensitivity of elongating hypocotyls to sucrose,
368 because PIFs are required for sucrose-induced hypocotyl elongation (Stewart et al., 2011; Stewart
369 Lilley et al., 2012). This might explain the PAR-independent reduction in sucrose-induced
370 hypocotyl elongation that occurred under long photoperiods (Fig. 2C). **In the future**, it will be
371 informative to resolve the relative contributions of these mechanisms to sucrose-induced hypocotyl

372 elongation, given that Tre6P can regulate expression of both PIFs and auxin signalling genes (Paul
373 et al., 2010). This could provide insights into the nature of the coupling of SnRK1-mediated sugar
374 signalling and growth regulation by PIFs (Paul et al., 2010; Stewart et al., 2011; Stewart Lilley et
375 al., 2012).

376 *Involvement of phytohormone signals in sucrose-induced hypocotyl elongation under light/dark*
377 *cycles*

378 Auxin, GA and brassinosteroids are reported to mediate sucrose-induced hypocotyl elongation,
379 with a role for auxin identified under light/dark cycles and roles for GA and brassinosteroids
380 identified under extended darkness (de Lucas et al., 2008; Zhang et al., 2010; Liu et al., 2011;
381 Stewart et al., 2011; Stewart Lilley et al., 2012; Zhang et al., 2015; Zhang et al., 2016). Consistent
382 with this, our data indicate that auxin signalling has a major role in sucrose-induced hypocotyl
383 elongation under light/dark cycles (Fig. 4A), with GA signalling also contributing to this process
384 (Fig. 5B, C). We suggest two possible reasons why paclobutrazol completely abolished sucrose-
385 induced hypocotyl elongation (Fig. 5A), whereas the *gai-1* mutant only led to partial inhibition of
386 this phenotype (Fig. 5B). One possibility is that DELLA-independent GA signalling contributes to
387 sucrose-induced hypocotyl elongation, since DELLA proteins control around 40-60% of GA-
388 regulated transcripts (Cao et al., 2006). An alternative possibility is that these were off-target or
389 ectopic effects of paclobutrazol, because the paclobutrazol-induced attenuation of hypocotyl
390 elongation was not rescued fully by GA supplementation (Fig. 5A).

391 Auxin-induced expansins that are upregulated during hypocotyl elongation were also induced by
392 sucrose supplementation (Fig. 4B-E; Fig. S2). Whilst *EXPA11* was induced strongly by sucrose,
393 the small response of *EXPA8* to sucrose in the wild type makes it difficult to interpret the responses
394 of *EXPA8* to sucrose in KIN10-ox and the *tps1* mutants (Fig. 4B, C). Interestingly, sucrose

induction of *EXPA11* was abolished in KIN10-ox, suggesting a role for KIN10 in expansin gene expression within elongating hypocotyls. In comparison, these expansins were sucrose-inducible in *tps1-11* and *tps1-12* (Fig. 4B-E). One possible explanation is that KIN10-ox causes a much greater level of SnRK1 activity compared with the *tps* mutants, which are hypomorphic alleles that harbour reduced Tre6P concentrations (Gómez et al., 2010) and are not completely deficient in sucrose-induced hypocotyl elongation (Fig. 1D, E).

An alternative and speculative explanation for the different behaviour of expansin transcripts in KIN10-ox and *tps* mutants could relate to Tre6P-KIN10 regulating growth through two broad processes- firstly, through direct signalling effects upon growth (e.g. by regulating auxin signals), and secondly through metabolic effects, such as growth constraints due to altered nocturnal catabolism. This could point to TPS1 and SnRK1 making independent contributions to sucrose-induced hypocotyl elongation under light/dark cycles, potentially through separate signalling and metabolic effects, rather than acting in series. Our data suggest that sucrose-induced hypocotyl elongation under light/dark cycles includes a signalling effect, previously proposed to occur through PIF-regulated auxin signals (Stewart et al., 2011; Stewart Lilley et al., 2012). On the other hand, the unexpected behaviour of expansin transcripts in *tps1* mutants (Fig. 1D, E) suggests that mechanisms additional to auxin/GA signalling might contribute to sucrose-induced hypocotyl elongation under light/dark cycles. These additional mechanisms could involve brassinosteroid and/or TOR signalling, which are required for sucrose-induced increases in hypocotyl length under extended darkness (Zhang et al., 2015; Zhang et al., 2016). It would be informative in future to investigate the crosstalk between SnRK1 and TOR energy signalling during hypocotyl elongation, to gain insights into the relative importance of these energy management pathways to the below-ground (darkness) and above-ground (light/dark cycles) stages of seedling establishment.

Conclusions

419 We identified a novel role for the SnRK1 energy signalling hub in the regulation of sucrose-
420 induced hypocotyl elongation under light/dark cycles. We propose that KIN10 could be positioned
421 upstream from the auxin and GA signals that lead to sucrose-induced hypocotyl elongation in the
422 light (Liu et al., 2011; Stewart et al., 2011; Stewart Lilley et al., 2012). A question for future
423 investigation concerns the functional organization of this pathway. In one scenario, KIN10-
424 mediated energy signalling regulates hypocotyl elongation by acting upon phytohormone
425 signalling, potentially through PIFs (Stewart Lilley *et al.*, 2012). In a different and non-exclusive
426 scenario, SnRK1-mediated alterations in metabolic enzyme activity and growth-related transcripts
427 prime hypocotyls to capitalize upon increased sucrose availability (Nunes *et al.*, 2013a). This is an
428 interesting question in the case of hypocotyl elongation, which arises from cell expansion rather
429 than growth through cell division and biomass accumulation *per se* (Gendreau *et al.*, 1997). These
430 two possibilities are non-exclusive, because the phenotypic differences that we report between
431 KIN10-ox lines and *tps1* mutants (e.g. expansin transcript accumulation; Fig. 4) could implicate
432 more than one mechanism in sucrose-induced hypocotyl elongation.

433 A further question for future investigation is of the nature of the interplay between KIN10/Tre6P,
434 TOR and brassinosteroids in the regulation of hypocotyl elongation in response to sugars. One
435 speculative hypothesis is that under conditions of starvation, such as when a developing below-
436 ground seedling is exhausting its seed-based energy store, brassinosteroid signalling produces a
437 strong elongation cue to drive seedling emergence into the light (Zhang et al., 2015; Zhang et al.,
438 2016). Then, once the seedling has emerged into the daily cycles of light and dark, KIN10/Tre6P
439 adjusts the elongation of hypocotyls to allow optimal seedling establishment under local light
440 conditions (Fig. 1, Fig. 2). It is possible that increased SnRK1 activity under conditions of
441 transiently low light, for example due to unpredictable changes in the weather, operates alongside
442 phototransduction pathways to prevent inappropriate etiolation following seedling emergence.

443 Therefore, one potential function of the mechanism that we identified might be to adapt the rate of
444 seedling development to optimize the use of seed and photosynthetic resources under fluctuating
445 light environments.

446

447 **Materials and Methods**

448 *Plant material and growth conditions*

449 *Arabidopsis* (*Arabidopsis thaliana* (L.) Heynh.) seeds were surface-sterilized and sown on half-
450 strength Murashige & Skoog basal salt mixture (Duchefa, Netherlands) (0.5 MS) with 0.8% (w/v)
451 agar (Noordally et al., 2013). Seeds were then stratified (3 days at 4 °C) and germinated and grown
452 for 7 days under 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of white light at 19 °C, except Fig. 2B-E where PAR was
453 reduced. Media was supplemented with either 3 % (w/v) sucrose (87.6 mM) or 87.6 mM sorbitol as
454 an osmotic control, according to the experiment. For experiments investigating gibberellin
455 signalling, media was supplemented with 20 μM paclobutrazol (PAC) and 100 μM gibberellic acid
456 (GA_3 form) (both Sigma-Aldrich) with a methanol carrier. Paclobutrazol is effective for studies of
457 GA signalling during development at the concentration of 20 μM (Penfield et al., 2004; MacGregor
458 et al., 2015). For experiments investigating auxin signalling, media was supplemented with 1-N-
459 naphthylphthalamic acid (NPA, Sigma-Aldrich) at up to 10 μM with a dimethylsulfoxide (DMSO)
460 carrier. Controls were supplemented with the appropriate carrier at the same concentration as
461 treatment media (0.1% (v/v) DMSO for NPA; 0.12% (v/v) methanol for PAC and GA).
462 To transfer growing seedlings to media containing GA or PAC, surface sterilized and stratified
463 seeds were pipetted onto 1 μm pore-diameter nylon mesh (Normesh, UK), on top of 0.5 MS 0.8%
464 (w/v) agar, and allowed to germinate for 3 days. Seedlings were then transferred to 0.5 MS
465 supplemented with either 3% (w/v) sucrose (87.6 mM) or 87.6 mM sorbitol, plus 20 μM PAC, 100

466 μ M GA or both PAC and GA. Hypocotyls were measured after 5 days growth on treatment plates.
467 For experiments with circadian oscillator mutants, we did not use arrhythmic CCA1-ox plants
468 because overexpression of CCA1 causes very long hypocotyls (Wang and Tobin, 1998), which
469 would confound investigation of the role of sugars in hypocotyl elongation.
470 Genotypes used were *tps1* TILLING mutants (Gómez et al., 2010), KIN10-ox (Baena-González et
471 al., 2007), *gin2-1* (Moore et al., 2003), *gai-1* (Koorneef et al., 1985), DELLA global mutant (Koini
472 et al., 2009), *pyr1 pyl1 pyl2 pyl4* (Park et al., 2009), *cca1-11 lhy-21 toc1-21* (Ding et al., 2007), *gi-*
473 *11* (Richardson et al., 1998) and *prp7-11* (Yamamoto et al., 2003; Nakamichi et al., 2005). In the
474 KIN10-ox lines, *KIN10* transcript abundance was 17-fold greater than the wild type in elongating
475 hypocotyls (Fig. S6A). In the *tps1-11* and *tps1-12* alleles, *TPS1* transcript abundance was
476 unchanged (*tps1-11*) or slightly increased (*tps1-12*) compared with the wild type (Fig. S6B). This
477 result for the *tps1* alleles was unsurprising because these are mis-sense mutants rather than
478 insertion mutants (Gómez et al., 2010).

479 *Hypocotyl measurement*

480 Seedlings were grown on square petri dishes within temperature-controlled growth chambers
481 (Panasonic MLR-352). Plates were angled at about 45 degrees to allow hypocotyls to elongate
482 without touching lids. Hypocotyls were measured by positioning 7 day-old seedlings on the surface
483 of 1% (w/v) agar for photography (Nikon D50) and subsequent measurement using the ImageJ
484 software (<https://imagej.nih.gov/ij/>).

485 *RNA extraction and qRT-PCR*

486 RNA was extracted according to (Noordally et al., 2013), using the Machery-Nagel Nucleospin II
487 plant RNA extraction kit incorporating DNase I treatment (Thermo-Fisher), except approximately
488 60 seedlings were used per RNA sample. cDNA was synthesized using the High Capacity cDNA

Reverse Transcription Kit with RNase Inhibitor (Applied Biosystems), according to manufacturer's instructions. cDNA was analyzed using an MXPro 3005 real time PCR system (Agilent) with Brilliant III Ultra-Fast SYBR qPCR mastermix (Agilent) (primers in Table S1). At least two technical repeats were performed for each qRT-PCR reaction. Data were analyzed using the $\Delta\Delta C_t$ method, with *PROTEIN PHOSPHATASE 2A SUBUNIT A3 (PP2AA3)* as a reference transcript.

494

Supplemental Material

Figure S1. The *cca1-11 lhy-21 toc1-21* triple mutant does not alter sucrose-induced hypocotyl elongation (direct repeat of Figure 3A).

Figure S2. Selection of expansin transcripts for experimentation.

Figure S3. Sucrose supplementation of growth media did not alter abundance of auxin biosynthesis transcripts or auxin-responsive transcripts relative to osmotic controls.

Figure S4. Efficacy of GA₃ used for study.

Figure S5. ABA signalling is not required for sucrose-induced hypocotyl elongation under short photoperiods.

Figure S6. *KIN10* and *TPS1* transcript abundance in KIN10-ox and *tps1* TILLING mutants.

Table S1. qRT-PCR primer sequences.

506

507

Acknowledgements

We thank Ian Graham (York), Filip Rolland (KU Leuven), Kerry Franklin (Bristol), Jean-Charles Isner (Bristol), Nicholas Harberd (Oxford) and Alex Webb (Cambridge) for seed donation, and Ian Graham for the discussion and encouragement that led to this work. We thank Kerry Franklin and

512 Brendan Davies (Leeds) for critical feedback. This research was funded by the BBSRC (UK)
513 (grant BB/I005811/2; South-West Doctoral Training Partnership BB/J014400/1), the Lady Emily
514 Smyth Agricultural Research Station (Bristol), the Wolfson Foundation and The Royal Society.
515 Antony Dodd is grateful to Kyoto University for awarding a Guest Professorship (Joint
516 Usage/Research Program of the Center for Ecological Research, Kyoto University) and to The
517 Royal Society for awarding a University Research Fellowship.

518

519 **Figure legends**

520 **Figure 1.** KIN10 and TPS1 participate in sucrose-induced hypocotyl elongation. (A)

521 Representative images of *L. er.* wild type, KIN10-ox and *tps1* seedlings cultivated under a variety
522 of photoperiods, with and without supplementation with 3% sucrose. All panels scaled identically.

523 Images are a subset of seedlings used to generate data in (B-E). (B-E) Lengths of hypocotyls of
524 seedlings grown under (B, D) constant light, 16 h and 8 h photoperiods, and (C, E) 4 h

525 photoperiods. Photoperiods are indicated underneath graphs. (F) Effect of sucrose supplementation
526 upon *gin2-1* hypocotyl length. S.E.M. is small under continuous light (0.03 – 0.05 mm), so not
527 visible on graphs. Data were analysed with ANOVA and Tukey's post-hoc tests ($n = 10$ (B-E) or n
528 $= 20$ (F) seedlings in three independent experiments, \pm S.E.M). Different letters indicate

529 statistically significant differences between means, specifically within each light condition ($p <$
530 0.05). (B-E); MS is half-strength MS media, and Suc and Sor are 0.5 MS supplemented with 3%
531 (w/v) sucrose or equimolar sorbitol (87.6 mM osmotic control), respectively.

532

533 **Figure 2.** Day-length dependency of sucrose-induced hypocotyl elongation in wild type seedlings.

534 (A) Increase in hypocotyl length caused by sucrose under range of photoperiods (data derived from

535 Fig. 1, plotted relative to sorbitol control). (B-E) Comparison of (B, D) absolute hypocotyl length

536 and (C, E) proportional increase in hypocotyl length caused by sucrose supplementation under
 537 specified photosynthetically active radiation (PAR) and photoperiod. Mean \pm S.E.M; (A, C-E)
 538 $n = 10$ seedlings in two independent experiments (B) $n = 20$ seedlings. Data analysed using
 539 ANOVA followed by post-hoc Tukey test. Different letters indicate statistically significant
 540 differences between means ($p < 0.05$).

541

542 **Figure 3.** The circadian oscillator does not participate in sucrose-induced hypocotyl elongation
 543 under short photoperiods. Sucrose-induced change in hypocotyl length of (A) a circadian oscillator
 544 triple mutant (*cca1-11 lhy-21 toc1-21*, background Ws-2) and (B, C) two oscillator components
 545 participating in sucrose regulation of the circadian oscillator. (D) Change in hypocotyl length
 546 caused by sucrose supplementation in *gi-11* and *prp7-11*, expressed relative to 0.5 MS control. MS
 547 is 0.5 MS media, and Suc and Sor are 0.5 MS supplemented with 3% (w/v) sucrose and sorbitol
 548 (87.6 mM, osmotic control), respectively. Data are mean \pm S.E.M ($n = 10 - 16$), analysed with (A-
 549 C) ANOVA and post-hoc Tukey tests and (D) two-sample t-test comparing mutant with wild type
 550 for each treatment. Data show one of three independent repeats of the experiment, conducted under
 551 4 h photoperiods. Different letters indicate statistically significant differences between means ($p <$
 552 0.05).

553

554 **Figure 4.** Auxin signalling underlies sucrose-induced hypocotyl elongation and KIN10 regulates
 555 expansin gene expression. (A) Hypocotyl length of seedlings cultivated with a range of
 556 concentrations of the inhibitor of polar auxin transport 1-N-naphthylphthalamic acid (NPA), under
 557 4 h photoperiods (mean \pm S.E.M; $n = 20$). (B-E) Sucrose-induced changes in expansin transcript
 558 abundance in elongating wild type, *tps1* and KIN10-ox seedlings under 4 h photoperiods. (B, D)
 559 Indicate *EXPA8* and *EXPA11* transcript abundance relative to *PP2AA3* (mean \pm S.E.M; $n = 3$). (C,

560 E) Indicate the magnitude of sucrose-induced change in transcript abundance in each genotype
561 relative to the osmotic control. Data analysed with ANOVA and post-hoc Tukey tests, and with
562 statistical significance indicated using starring (N.S. = not significant $p > 0.05$; * = $p \leq 0.05$; ** =
563 $p < 0.01$; *** = $p < 0.001$).

564

565 **Figure 5.** Gibberellin signals contribute to sucrose-induced hypocotyl elongation under short
566 photoperiods. (A) The GA biosynthesis inhibitor paclobutrazol (PAC) at 20 μ M inhibits sucrose-
567 induced hypocotyl elongation. Seedlings were germinated on MS agar and transferred to treatment
568 media after germination; carrier control was 0.12% (v/v) methanol. (B) Sucrose-induced hypocotyl
569 elongation was attenuated in *gai-1* mutant seedlings. (C) Sucrose-induced hypocotyl elongation
570 was unaltered in a DELLA global knockout mutant. Experiments performed under 4 h
571 photoperiods. Data are mean \pm S.E.M ($n = 20$) from one of two independent repeats, analysed with
572 ANOVA and post-hoc Tukey tests. Different letters indicate statistically significant differences
573 between means ($p < 0.05$). Osmotic control was 87.6 mM sorbitol.

574

575 **Supplemental Figure Legends**

576

577 **Figure S1.** The *cca1-11 lhy-21 toc1-21* triple mutant does not alter sucrose-induced hypocotyl
578 elongation under light/dark cycles. This is a direct repeat of the experiment in Figure 3A where
579 data approach statistical significance. (A) Comparison of hypocotyl length of Ws-2 background
580 and *cca1-11 lhy-21 toc1-21* grown on 0.5 MS media (MS) and 0.5 MS media supplemented with
581 3% (w/v) sucrose (Suc); (B) Increase in hypocotyl length of wild type and *cca1-11 lhy-21 toc1-21*
582 caused by exogenous sucrose, relative to 0.5 MS control. Data are mean \pm S.E.M; $n = 10$; statistical

583 significance from two-sample t-tests comparing mutant and wild type for each treatment; N.S. = no
584 significant difference ($p \geq 0.05$).

585

586 **Figure S2.** *EXPA8* and *EXPA11* transcripts were (A) up-regulated by conditions that promote
587 hypocotyl elongation (constant darkness) and (B) down-regulated by the auxin transport inhibitor
588 NPA (mean \pm S.E.M.; $n = 3$). Transcript abundance was relative to *PP2AA3* reference transcript
589 and used 7-day old *L. er.* seedlings. Data analysed with ANOVA followed by post-hoc Tukey test.
590 Different letters indicate statistically significant differences between means ($p < 0.05$).

591

592 **Figure S3.** Sucrose supplementation did not alter the abundance of auxin biosynthesis transcripts
593 or auxin-responsive transcripts relative to osmotic controls, due to responses of osmotic controls.
594 Data indicate relative abundance of three auxin biosynthesis transcripts (*YUCCA8*, *YUCCA9*,
595 *CYP79B3*) and two auxin-responsive transcripts (*IAA29*, *SAUR15*) in two backgrounds, using
596 *PP2AA3* as the reference transcript. Seedlings (60 per replicate) were grown on 0.5 MS, 3% (w/v)
597 sucrose, or 87.6 mM sorbitol as osmotic control, and harvested for RNA 4 days and 7 days after
598 germination (indicated on x axis). Two background lines were used to evaluate whether there were
599 ecotype-specific phenotypes. Data are mean \pm S.E.M; $n = 2$ independent biological repeats.
600 Analyzed by ANOVA ($p \geq 0.05$ in all cases, i.e. not significant).

601

602 **Figure S4.** Confirmation of activity of GA_3 . 100 μ M GA_3 increased hypocotyl length relative to
603 the carrier control in both *L. er.* and Col-0 backgrounds, under 4 h and 16 h photoperiods.
604 Seedlings were germinated and grown in presence of GA . Data were collected during methods
605 development and are not directly comparable with other experiments. Data expressed as mean \pm

606 S.E.M. ($n = 20$) and analysed with ANOVA followed by post-hoc Tukey test. Different letters
607 indicate statistically significant differences between means ($p < 0.05$).

608

609 **Figure S5.** ABA signalling is not required for sucrose-induced hypocotyl elongation under short
610 photoperiods. The *pyr1-1 pyl1-1 pyl2-1 pyl4-1* quadruple mutant incorporates Col-0 and *L. er.*
611 backgrounds (Park et al., 2009), both of which are included as controls. Data indicate mean
612 hypocotyl lengths of seedlings grown on 0.5 MS supplemented with 3% sucrose or an osmotic
613 control (87.6 mM sorbitol), under 4 h photoperiods. Data are mean \pm S.E.M.; $n = 20$ (background
614 lines); $n = 3 - 9$ depending on treatment for *pyr1-1 pyl1-1 pyl2-1 pyl4-1* (low replicate numbers
615 due to poor mutant germination). Data are from one of two independent repeats. Statistical
616 significance from independent-samples Kruskal-Wallis analysis of variance on ranks and post-hoc
617 Dunn tests comparing mutant and wild type for each treatment; *** = $p < 0.001$; N.S. = no
618 significant difference ($p \geq 0.05$).

619

620 **Figure S6.** *KIN10* and *TPS1* transcript abundance KIN10-ox and *tps1* TILLING mutants. (A)
621 *KIN10* transcript abundance in two independent KIN10-ox lines (Baena-González et al., 2007), its
622 *L. er* background, and also Col-0. Transcript abundance is relative to *PP2AA3* reference. (B) *TPS1*
623 transcript abundance in *tps1-11* and *tps1-12* (Gómez et al., 2010), alongside the *L. er* and Col-0
624 backgrounds. Transcript abundance was measured in 7 day old seedlings and is relative to the
625 *PP2AA3* reference transcript. Data expressed as mean \pm S.E.M ($n = 3$) and analyzed with ANOVA
626 followed by post-hoc Tukey test. Different letters indicate statistically significant differences
627 between means ($p < 0.05$).

628

629 **References**

630
631
632 **Avonce N, Leyman B, Mascorro-Gallardo JO, Van Dijck P, Thevelein JM, Iturriaga G** (2004)
633 The Arabidopsis trehalose-6-P synthase *AtTPS1* gene is a regulator of glucose, abscisic
634 acid, and stress signaling. *Plant Physiology* **136**: 3649-3659
635 **Baena-González E, Rolland F, Thevelein JM, Sheen J** (2007) A central integrator of
636 transcription networks in plant stress and energy signalling. *Nature* **448**: 938-942
637 **Baena-González E, Sheen J** (2008) Convergent energy and stress signaling. *Trends in Plant*
638 *Science* **13**: 474-482
639 **Belin C, Megies C, Hauserová E, Lopez-Molina L** (2009) Absciscic acid represses growth of the
640 Arabidopsis embryonic Axis after germination by enhancing auxin signaling. *The Plant*
641 *Cell* **21**: 2253-2268
642 **Boylan MT, Quail PH** (1991) Phytochrome A overexpression inhibits hypocotyl elongation in
643 transgenic Arabidopsis. *Proceedings of the National Academy of Sciences* **88**: 10806-
644 10810
645 **Cao D, Cheng H, Wu W, Soo HM, Peng J** (2006) Gibberellin mobilizes distinct DELLA-
646 dependent transcriptomes to regulate seed germination and floral development in
647 Arabidopsis. *Plant Physiology* **142**: 509-525
648 **Casal JJ** (2013) Photoreceptor signaling networks in plant responses to shade. *Annual Review of*
649 *Plant Biology* **64**: 403-427
650 **Cheng H, Qin L, Lee S, Fu X, Richards DE, Cao D, Luo D, Harberd NP, Peng J** (2004)
651 Gibberellin regulates Arabidopsis floral development via suppression of DELLA protein
652 function. *Development* **131**: 1055-1064
653 **Collett CE, Harberd NP, Leyser O** (2000) Hormonal interactions in the control of Arabidopsis
654 hypocotyl elongation. *Plant Physiology* **124**: 553-562
655 **Cowling RJ, Harberd NP** (1999) Gibberellins control Arabidopsis hypocotyl growth via
656 regulation of cellular elongation. *Journal of Experimental Botany* **50**: 1351-1357
657 **Crawford AJ, McLachlan DH, Hetherington AM, Franklin KA** (2012) High temperature
658 exposure increases plant cooling capacity. *Current Biology* **22**: R396-R397
659 **Dalchau N, Baek SJ, Briggs HM, Robertson FC, Dodd AN, Gardner MJ, Stancombe MA,**
660 **Haydon MJ, Stan G-B, Gonçalves JM, Webb AAR** (2011) The circadian oscillator gene
661 *GIGANTEA* mediates a long-term response of the Arabidopsis thaliana circadian clock to
662 sucrose. *Proceedings of the National Academy of Sciences* **108**: 5104-5109
663 **de Lucas M, Daviere JM, Rodriguez-Falcon M, Pontin M, Iglesias-Pedraz JM, Lorain S,**
664 **Fankhauser C, Blazquez MA, Titarenko E, Prat S** (2008) A molecular framework for
665 light and gibberellin control of cell elongation. *Nature* **451**: 480-484
666 **Debast S, Nunes-Nesi A, Hajirezaei MR, Hofmann J, Sonnewald U, Fernie AR, Börnke F**
667 (2011) Altering trehalose-6-phosphate content in transgenic potato tubers affects tuber
668 growth and alters responsiveness to hormones during sprouting. *Plant Physiology* **156**:
669 1754-1771
670 **Delatte TL, Sedijani P, Kondou Y, Matsui M, de Jong GJ, Somsen GW, Wiese-Klinkenberg**
671 **A, Primavesi LF, Paul MJ, Schluepmann H** (2011) Growth arrest by trehalose-6-
672 phosphate: An astonishing case of primary metabolite control over growth by way of the
673 SnRK1 signaling pathway. *Plant Physiology* **157**: 160
674 **Delorge I, Figueroa CM, Feil R, Lunn JE, van Dijck P** (2015) Trehalose-6-phosphate synthase 1
675 is not the only active TPS in *Arabidopsis thaliana*. *Biochemical Journal* **466**: 283-290

676 **Ding Z, Doyle MR, Amasino RM, Davis SJ** (2007) A complex genetic interaction between
677 *Arabidopsis thaliana* TOC1 and CCA1/LHY in driving the circadian clock and in output
678 regulation. *Genetics* **176**: 1501-1510

679 **Dobrenel T, Marchive C, Sormani R, Moreau M, Mozzo M, Montané M-H, Menand B,**
680 **Robaglia C, Meyer C** (2011) Regulation of plant growth and metabolism by the TOR
681 kinase. *Biochemical Society Transactions* **39**: 477

682 **Dowson-Day MJ, Millar AJ** (1999) Circadian dysfunction causes aberrant hypocotyl elongation
683 patterns in *Arabidopsis*. *The Plant Journal* **17**: 63-71

684 **Esmon CA, Tinsley AG, Ljung K, Sandberg G, Hearne LB, Liscum E** (2006) A gradient of
685 auxin and auxin-dependent transcription precedes tropic growth responses. *Proceedings of*
686 *the National Academy of Sciences of the United States of America* **103**: 236-241

687 **Fu X, Richards DE, Ait-ali T, Hynes LW, Ougham H, Peng J, Harberd NP** (2002) Gibberellin-
688 mediated proteasome-dependent degradation of the barley DELLA protein SLN1 repressor.
689 *The Plant Cell* **14**: 3191-3200

690 **Gangappa SN, Kumar SV** (2017) DET1 and HY5 control PIF4-mediated thermosensory
691 elongation growth through distinct mechanisms. *Cell Reports* **18**: 344-351

692 **Gendreau E, Traas J, Desnos T, Grandjean O, Caboche M, Hofte H** (1997) Cellular basis of
693 hypocotyl growth in *Arabidopsis thaliana*. *Plant Physiology* **114**: 295-305

694 **Goda H, Sawa S, Asami T, Fujioka S, Shimada Y, Yoshida S** (2004) Comprehensive
695 comparison of auxin-regulated and brassinosteroid-regulated genes in *Arabidopsis*. *Plant*
696 *Physiology* **134**: 1555-1573

697 **Gómez LD, Baud S, Gilday A, Li Y, Graham IA** (2006) Delayed embryo development in the
698 *ARABIDOPSIS* TREHALOSE-6-PHOSPHATE SYNTHASE 1 mutant is associated with
699 altered cell wall structure, decreased cell division and starch accumulation. *The Plant*
700 *Journal* **46**: 69-84

701 **Gómez LD, Gilday A, Feil R, Lunn JE, Graham IA** (2010) AtTPS1-mediated trehalose 6-
702 phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness
703 to ABA in germinating seeds and stomatal guard cells. *The Plant Journal* **64**: 1-13

704 **Graf A, Schlereth A, Stitt M, Smith AM** (2010) Circadian control of carbohydrate availability
705 for growth in *Arabidopsis* plants at night. *Proceedings of the National Academy of Sciences*
706 **107**: 9458-9463

707 **Griffiths CA, Sagar R, Geng Y, Primavesi LF, Patel MK, Passarelli MK, Gilmore IS, Steven**
708 **RT, Bunch J, Paul MJ, Davis BG** (2016) Chemical intervention in plant sugar signalling
709 increases yield and resilience. *Nature* **540**: 574-578

710 **Haydon MJ, Mielczarek O, Robertson FC, Hubbard KE, Webb AAR** (2013) Photosynthetic
711 entrainment of the *Arabidopsis thaliana* circadian clock. *Nature* **502**: 689-692

712 **Hayes S, Velanis CN, Jenkins GI, Franklin KA** (2014) UV-B detected by the UVR8
713 photoreceptor antagonizes auxin signaling and plant shade avoidance. *Proceedings of the*
714 *National Academy of Sciences* **111**: 11894-11899

715 **Jang JC, León P, Zhou L, Sheen J** (1997) Hexokinase as a sugar sensor in higher plants. *The*
716 *Plant Cell* **9**: 5-19

717 **Keuskamp DH, Pollmann S, Voesenek LACJ, Peeters AJM, Pierik R** (2010) Auxin transport
718 through PIN-FORMED 3 (PIN3) controls shade avoidance and fitness during competition.
719 *Proceedings of the National Academy of Sciences* **107**: 22740-22744

720 **Kim BC, Tennessen DJ, Last Robert L** (1998) UV-B-induced photomorphogenesis in
721 *Arabidopsis thaliana*. *The Plant Journal* **15**: 667-674

- Koini MA, Alvey L, Allen T, Tilley CA, Harberd NP, Whitelam GC, Franklin KA** (2009) High temperature-mediated adaptations in plant architecture require the bHLH transcription factor PIF4. *Current Biology* **19**: 408-413
- Koorneef M, Elgersma A, Hanhart CJ, van Loenen-Martinet EP, van Rijn L, Zeevaart JAD** (1985) A gibberellin insensitive mutant of *Arabidopsis thaliana*. *Physiologia Plantarum* **65**: 33-39
- Kurata T, Yamamoto KT** (1998) *petit1*, a conditional growth mutant of *Arabidopsis* defective in sucrose-dependent elongation growth. *Plant Physiology* **118**: 793
- Lee DJ, Park JW, Lee HW, Kim J** (2009) Genome-wide analysis of the auxin-responsive transcriptome downstream of *iaa1* and its expression analysis reveal the diversity and complexity of auxin-regulated gene expression. *Journal of Experimental Botany* **60**: 3935-3957
- Leivar P, Quail PH** (2011) PIFs: pivotal components in a cellular signaling hub. *Trends in Plant Science* **16**: 19-28
- Li Y, Darley CP, Ongaro V, Fleming A, Schipper O, Baldauf SL, McQueen-Mason SJ** (2002) Plant expansins are a complex multigene family with an ancient evolutionary origin. *Plant Physiology* **128**: 854-864
- Li Y, Van den Ende W, Rolland F** (2014) Sucrose induction of anthocyanin biosynthesis is mediated by DELLA. *Molecular Plant* **7**: 570-572
- Lincoln C, Britton JH, Estelle M** (1990) Growth and development of the *axr1* mutants of *Arabidopsis*. *Plant Cell* **2**: 1071-1080
- Liscum E, Hangarter RP** (1991) *Arabidopsis* mutants lacking blue light-dependent inhibition of hypocotyl elongation. *The Plant Cell* **3**: 685-694
- Liu Z, Zhang Y, Liu R, Hao H, Wang Z, Bi Y** (2011) Phytochrome interacting factors (PIFs) are essential regulators for sucrose-induced hypocotyl elongation in *Arabidopsis*. *Journal of Plant Physiology* **168**: 1771-1779
- Lorrain S, Allen T, Duek PD, Whitelam GC, Fankhauser C** (2008) Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *The Plant Journal* **53**: 312-323
- Lunn JE, Feil R, Hendriks JHM, Gibon Y, Morcuende R, Osuna D, Sheible WR, Carillo P, Hajirezaei MR, Stitt M** (2006) Sugar-induced increases in trehalose-6-phosphate are correlated with redox activation of ADPglucose pyrophosphorylase and higher rates of starch synthesis in *Arabidopsis thaliana*. *Biochemical Journal* **397**: 139-148
- MacGregor DR, Kendall SL, Florance H, Fedi F, Moore K, Paszkiewicz K, Smirnoff N, Penfield S** (2015) Seed production temperature regulation of primary dormancy occurs through control of seed coat phenylpropanoid metabolism. *New Phytologist* **205**: 642-652
- Mair A, Pedrotti L, Wurzinger B, Anrather D, Simeunovic A, Weiste C, Valerio C, Dietrich K, Kirchler T, Nägele T, Vicente Carbajosa J, Hanson J, Baena-González E, Chaban C, Weckwerth W, Dröge-Laser W, Teige M** (2015) SnRK1-triggered switch of bZIP63 dimerization mediates the low-energy response in plants. *eLife* **4**
- Más P, Alabadí D, Yanovsky MJ, Oyama T, Kay SA** (2003) Dual role of TOC1 in the control of circadian and photomorphogenic responses in *Arabidopsis*. *The Plant Cell* **15**: 223-236
- Mattioli CC, Tomaz JP, Duarte GT, Prado FM, Del Bem LEV, Silveira AB, Gauer L, Corrêa LGG, Drumond RD, Viana AJC, Di Mascio P, Meyer C, Vincentz M** (2011) The *Arabidopsis* bZIP gene *AtbZIP63* is a sensitive integrator of transient abscisic acid and glucose signals. *Plant Physiology* **157**: 692-705

768 **Miyazaki Y, Jikumaru Y, Takase T, Saitoh A, Sugitani A, Kamiya Y, Kiyosue T** (2016)
769 Enhancement of hypocotyl elongation by LOV KELCH PROTEIN2 production is mediated
770 by auxin and phytochrome-interacting factors in *Arabidopsis thaliana*. *Plant Cell Reports*
771 **35**: 455-467

772 **Mizuno T, Nomoto Y, Oka H, Kitayama M, Takeuchi A, Tsubouchi M, Yamashino T** (2014)
773 Ambient temperature signal feeds into the circadian clock transcriptional circuitry through
774 the EC night-time repressor in *Arabidopsis thaliana*. *Plant and Cell Physiology* **55**: 958-976

775 **Moore B, Zhou L, Rolland F, Hall Q, Cheng W-H, Liu Y-X, Hwang I, Jones T, Sheen J**
776 (2003) Role of the *Arabidopsis* glucose sensor HXK1 in nutrient, light, and hormonal
777 signaling. *Science* **300**: 332-336

778 **Nakamichi N, Kita M, Ito S, Yamashino T, Mizuno T** (2005) PSEUDO-RESPONSE
779 REGULATORS, PRR9, PRR7 and PRR5, together play essential roles close to the
780 circadian clock of *Arabidopsis thaliana*. *Plant and Cell Physiology* **46**: 686-698

781 **Noordally ZB, Ishii K, Atkins KA, Wetherill SJ, Kusakina J, Walton EJ, Kato M, Azuma M,**
782 **Tanaka K, Hanaoka M, Dodd AN** (2013) Circadian control of chloroplast transcription by
783 a nuclear-encoded timing signal. *Science* **339**: 1316-1319

784 **Nozue K, Covington MF, Duek PD, Lorrain S, Fankhauser C, Harmer SL, Maloof JN** (2007)
785 Rhythmic growth explained by coincidence between internal and external cues. *Nature* **448**:
786 358-361

787 **Nuccio ML, Wu J, Mowers R, Zhou H-P, Meghji M, Primavesi LF, Paul MJ, Chen X, Gao Y,**
788 **Haque E, Basu SS, Lagrimini LM** (2015) Expression of trehalose-6-phosphate
789 phosphatase in maize ears improves yield in well-watered and drought conditions. *Nat*
790 *Biotech* **33**: 862-869

791 **Nunes C, O'Hara LE, Primavesi LF, Delatte TL, Schluepmann H, Somsen GW, Silva AB,**
792 **Feverieiro PS, Wingler A, Paul MJ** (2013) The trehalose 6-phosphate/SnRK1 signaling
793 pathway primes growth recovery following relief of sink limitation. *Plant Physiology* **162**:
794 1720-1732

795 **Nunes C, Primavesi LF, Patel MK, Martinez-Barajas E, Powers SJ, Sagar R, Feverieiro PS,**
796 **Davis BG, Paul MJ** (2013) Inhibition of SnRK1 by metabolites: Tissue-dependent effects
797 and cooperative inhibition by glucose 1-phosphate in combination with trehalose 6-
798 phosphate. *Plant Physiology and Biochemistry* **63**: 89-98

799 **Nusinow DA, Helfer A, Hamilton EE, King JJ, Imaizumi T, Schultz TF, Farre EM, Kay SA**
800 (2011) The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of
801 hypocotyl growth. *Nature* **475**: 398-402

802 **Park S, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J, Rodrigues**
803 **A, Chow TF, Alfred SE, Bonetta D, Finkelstein R, Provart NJ, Desveaux D, Rodriguez**
804 **PL, McCourt P, Zhu JK, Schroeder JI, Volkman BF, Cutler SR** (2009) Absciscic acid
805 inhibits PP2Cs via the PYR/PYL family of ABA-binding START proteins. *Science* **324**:
806 1068-1071

807 **Paul MJ, Jhurrea D, Zhang Y, Primavesi LF, Delatte T, Schluepmann H, Wingler A** (2010)
808 Up-regulation of biosynthetic processes associated with growth by trehalose 6-phosphate.
809 *Plant Signaling & Behavior* **5**: 386-392

810 **Penfield S, Rylott EL, Gilday AD, Graham S, Larson TR, Graham IA** (2004) Reserve
811 mobilization in the *Arabidopsis* endosperm fuels hypocotyl elongation in the dark, is
812 independent of abscisic acid, and requires PHOSPHOENOLPYRUVATE
813 CARBOXYKINASE1. *The Plant Cell* **16**: 2705-2718

- Peng J, Carol P, Richards DE, King KE, Cowling RJ, Murphy GP, Harberd NP** (1997) The *Arabidopsis GAI* gene defines a signaling pathway that negatively regulates gibberellin responses *Genes & Development* **11**: 3194-3205
- Ramon M, Rolland F, Thevelein JM, van Dijck P** (2007) ABI4 mediates the effects of exogenous trehalose on *Arabidopsis* growth and starch breakdown. *Plant Molecular Biology* **63**: 195-206
- Richardson K, Fowler S, Pullen C, Skelton C, Morris B, Putterill J** (1998) T-DNA tagging of a flowering-time gene and improved gene transfer by *in planta* transformation of *Arabidopsis*. *Functional Plant Biology* **25**: 125-130
- Salter MG, Franklin KA, Whitelam GC** (2003) Gating of the rapid shade-avoidance response by the circadian clock in plants. *Nature* **426**: 680-683
- Schluepmann H, Pellny T, van Dijken A, Smeekens S, Paul M** (2003) Trehalose 6-phosphate is indispensable for carbohydrate utilization and growth in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the United States of America* **100**: 6849-6854
- Shin J, Sánchez-Villarreal A, Davis AM, Du S-x, Berendzen KW, Koncz C, Ding Z, Li C, Davis SJ** (2017) The metabolic sensor AKIN10 modulates the *Arabidopsis* circadian clock in a light-dependent manner. *Plant, Cell & Environment* **40**: 997-1008
- Smeekens S, Ma J, Hanson J, Rolland F** (2010) Sugar signals and molecular networks controlling plant growth. *Current Opinion in Plant Biology* **13**: 273-278
- Stewart JL, Maloof JN, Nemhauser JL** (2011) PIF genes mediate the effect of sucrose on seedling growth dynamics. *PLoS One* **6**: e19894
- Stewart Lilley JL, Gee CW, Sairanen I, Ljung K, Nemhauser JL** (2012) An endogenous carbon-sensing pathway triggers increased auxin flux and hypocotyl elongation. *Plant Physiology* **160**: 2261-2270
- Sulpice R, Flis A, Ivakov AA, Apelt F, Krohn N, Encke B, Abel C, Feil R, Lunn JE, Stitt M** (2014) *Arabidopsis* coordinates the diurnal regulation of carbon allocation and growth across a wide range of photoperiods. *Molecular Plant* **7**: 137-155
- Takahashi F, Sato-Nara K, Kobayashi K, Suzuki M, Suzuki H** (2003) Sugar-induced adventitious roots in *Arabidopsis* seedlings. *Journal of Plant Research* **116**: 83-91
- Vandesteene L, Ramon M, Le Roy K, Van Dijck P, Rolland F** (2010) A single active trehalose-6-P synthase (TPS) and a family of putative regulatory TPS-like proteins in *Arabidopsis*. *Molecular Plant* **3**: 406-419
- Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M** (2013) Regulation of flowering by trehalose-6-phosphate signaling in *Arabidopsis thaliana*. *Science* **339**: 704-707
- Wang Z-Y, Tobin EM** (1998) Constitutive expression of the *CIRCADIAN CLOCK ASSOCIATED 1 (CCA1)* gene disrupts circadian rhythms and suppresses its own expression. *Cell* **93**: 1207-1217
- Weinig C, Johnston JA, Willis CG, Maloof JN** (2007) Antagonistic multilevel selection on size and architecture in variable density settings. *Evolution* **61**: 58-67
- Wigge PA** (2013) Ambient temperature signalling in plants. *Current Opinion in Plant Biology* **16**: 661-666
- Wingler A, Delatte TL, O'Hara LE, Primavesi LF, Jhurreea D, Paul MJ, Schluepmann H** (2012) Trehalose 6-phosphate is required for the onset of leaf senescence associated with high carbon availability. *Plant Physiology* **158**: 1241-1251

- Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ** (2007) An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets. *PLoS One* **2**: e718
- Yadav UP, Ivakov A, Feil R, Duan GY, Walther D, Giavalisco P, Piques M, Carillo P, Hubberten H-M, Stitt M, Lunn JE** (2014) The sucrose–trehalose 6-phosphate (Tre6P) nexus: specificity and mechanisms of sucrose signalling by Tre6P. *Journal of Experimental Botany* **65**: 1051-1068
- Yamamoto Y, Sato E, Shimizu T, Nakamich N, Sato S, Kato T, Tabata S, Nagatani A, Yamashino T, Mizuno T** (2003) Comparative genetic studies on the *APRR5* and *APRR7* genes belonging to the *APRR1/TOC1* quintet implicated in circadian rhythm, control of flowering time, and early photomorphogenesis. *Plant and Cell Physiology* **44**: 1119-1130
- Zhang Y, Liu Z, Wang J, Chen Y, Bi Y, He J** (2015) Brassinosteroid is required for sugar promotion of hypocotyl elongation in *Arabidopsis* in darkness. *Planta* **242**: 881-893
- Zhang Y, Liu Z, Wang L, Zheng S, Xie J, Bi Y** (2010) Sucrose-induced hypocotyl elongation of *Arabidopsis* seedlings in darkness depends on the presence of gibberellins. *Journal of Plant Physiology* **167**: 1130-1136
- Zhang Y, Primavesi LF, Jhurreea D, Andralojc PJ, Mitchell RAC, Powers SJ, Schluepmann H, Delatte T, Wingler A, Paul MJ** (2009) Inhibition of SNF1-related protein kinase 1 activity and regulation of metabolic pathways by trehalose-6-phosphate. *Plant Physiology* **149**: 1860-1871
- Zhang Y, Shewry PR, Jones H, Barcelo P, Lazzeri PA, Halford NG** (2001) Expression of antisense SnRK1 protein kinase sequence causes abnormal pollen development and male sterility in transgenic barley. *The Plant Journal* **28**: 431-441
- Zhang Z, Zhu J-Y, Roh J, Marchive C, Kim S-K, Meyer C, Sun Y, Wang W, Wang Z-Y** (2016) TOR signaling promotes accumulation of BZR1 to balance growth with carbon availability in *Arabidopsis*. *Current Biology* **26**: 1854-1860




Figure 1. KIN10 and TPS1 participate in sucrose-induced hypocotyl elongation. (A) Representative images of *L. er.* wild type, KIN10-ox and *tps1* seedlings cultivated under a variety of photoperiods, with and without supplementation with 3% sucrose. All panels scaled identically. Images are a subset of seedlings used to generate data in (B-E). (B-E) Lengths of hypocotyls of seedlings grown under (B, D) constant light, 16 h and 8 h photoperiods, and (C, E) 4 h photoperiods. Photoperiods are indicated underneath graphs. (F) Effect of sucrose supplementation upon *gin2-1* hypocotyl length. S.E.M. is small under continuous light (0.03 – 0.05 mm), so not visible on graphs. Data were analysed with ANOVA and Tukey's post-hoc tests ($n = 10$ (B-E) or $n = 20$ (F) seedlings in three independent experiments, \pm S.E.M.). Different letters indicate statistically significant differences between means, specifically within each light condition ($p < 0.05$). (B-E); MS is half-strength MS media, and Suc and Sor are 0.5 MS supplemented with 3% (w/v) sucrose or equimolar sorbitol (87.6 mM osmotic control), respectively.




Figure 2. Day-length dependency of sucrose-induced hypocotyl elongation in wild type seedlings. (A) Increase in hypocotyl length caused by sucrose under range of photoperiods (data derived from Fig. 1, plotted relative to sorbitol control). (B-E) Comparison of (B, D) absolute hypocotyl length and (C, E) proportional increase in hypocotyl length caused by sucrose supplementation under specified photosynthetically active radiation (PAR) and photoperiod. Mean \pm S.E.M; (A, C-E) $n = 10$ seedlings in two independent experiments (B) $n = 20$ seedlings. Data analysed using ANOVA followed by post-hoc Tukey test. Different letters indicate statistically significant differences between means ($p < 0.05$).




Figure 3. The circadian oscillator does not participate in sucrose-induced hypocotyl elongation under short photoperiods. Sucrose-induced change in hypocotyl length of (A) a circadian oscillator triple mutant (*cca1-11 lhy-21 toc1-21*, background Ws-2) and (B, C) two oscillator components participating in sucrose regulation of the circadian oscillator. (D) Change in hypocotyl length caused by sucrose supplementation in *gi-11* and *prp7-11*, expressed relative to 0.5 MS control. MS is 0.5 MS media, and Suc and Sor are 0.5 MS supplemented with 3% (w/v) sucrose and sorbitol (87.6 mM, osmotic control), respectively. Data are mean \pm S.E.M ($n = 10 - 16$), analysed with (A-C) ANOVA and post-hoc Tukey tests and (D) two-sample t-test comparing mutant with wild type for each treatment. Data show one of three independent repeats of the experiment, conducted under 4 h photoperiods. Different letters indicate statistically significant differences between means ($p < 0.05$).




Figure 4. Auxin signalling underlies sucrose-induced hypocotyl elongation and KIN10 regulates expansin gene expression. (A) Hypocotyl length of seedlings cultivated with a range of concentrations of the inhibitor of polar auxin transport 1-N-naphthylphthalamic acid (NPA), under 4 h photoperiods (mean \pm S.E.M; $n = 20$). (B-E) Sucrose-induced changes in expansin transcript abundance in elongating wild type, *tps1* and KIN10-ox seedlings under 4 h photoperiods. (B, D) Indicate *EXPA8* and *EXPA11* transcript abundance relative to *PP2A43* (mean \pm S.E.M; $n = 3$). (C, E) Indicate the magnitude of sucrose-induced change in transcript abundance in each genotype relative to the osmotic control. Data analysed with ANOVA and post-hoc Tukey tests, and with statistical significance indicated using starring (N.S. = not significant $p > 0.05$; * = $p \leq 0.05$; ** = $p < 0.01$; *** = $p < 0.001$).




Figure 5. Gibberellin signals contribute to sucrose-induced hypocotyl elongation under short photoperiods. (A) The GA biosynthesis inhibitor paclobutrazol (PAC) at 20 μ M inhibits sucrose-induced hypocotyl elongation. Seedlings were germinated on MS agar and transferred to treatment media after germination; carrier control was 0.12% (v/v) methanol. (B) Sucrose-induced hypocotyl elongation was attenuated in *gai-1* mutant seedlings. (C) Sucrose-induced hypocotyl elongation was unaltered in a DELLA global knockout mutant. Experiments performed under 4 h photoperiods. Data are mean \pm S.E.M ($n = 20$) from one of two independent repeats, analysed with ANOVA and post-hoc Tukey tests. Different letters indicate statistically significant differences between means ($p < 0.05$). Osmotic control was 87.6 mM sorbitol.

Parsed Citations

Avonce N, Leyman B, Mascorro-Gallardo JO, Van Dijck P, Thevelein JM, Iturriaga G (2004) The Arabidopsis trehalose-6-P synthase AtTPS1 gene is a regulator of glucose, abscisic acid, and stress signaling. Plant Physiology 136: 3649-3659

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Baena-González E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of transcription networks in plant stress and energy signalling. Nature 448: 938-942

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Baena-González E, Sheen J (2008) Convergent energy and stress signaling. Trends in Plant Science 13: 474-482

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Belin C, Megies C, Hauserová E, Lopez-Molina L (2009) Absciscic acid represses growth of the Arabidopsis embryonic Axis after germination by enhancing auxin signaling. The Plant Cell 21: 2253-2268

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Boylan MT, Quail PH (1991) Phytochrome A overexpression inhibits hypocotyl elongation in transgenic Arabidopsis. Proceedings of the National Academy of Sciences 88: 10806-10810

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cao D, Cheng H, Wu W, Soo HM, Peng J (2006) Gibberellin mobilizes distinct DELLA-dependent transcriptomes to regulate seed germination and floral development in Arabidopsis. Plant Physiology 142: 509-525

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Casal JJ (2013) Photoreceptor signaling networks in plant responses to shade. Annual Review of Plant Biology 64: 403-427

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cheng H, Qin L, Lee S, Fu X, Richards DE, Cao D, Luo D, Harberd NP, Peng J (2004) Gibberellin regulates Arabidopsis floral development via suppression of DELLA protein function. Development 131: 1055-1064

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Collett CE, Harberd NP, Leyser O (2000) Hormonal interactions in the control of Arabidopsis hypocotyl elongation. Plant Physiology 124: 553-562

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cowling RJ, Harberd NP (1999) Gibberellins control Arabidopsis hypocotyl growth via regulation of cellular elongation. Journal of Experimental Botany 50: 1351-1357

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Crawford AJ, McLachlan DH, Hetherington AM, Franklin KA (2012) High temperature exposure increases plant cooling capacity. Current Biology 22: R396-R397

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dalchau N, Baek SJ, Briggs HM, Robertson FC, Dodd AN, Gardner MJ, Stancombe MA, Haydon MJ, Stan G-B, Gonçalves JM, Webb AAR (2011) The circadian oscillator gene GIGANTEA mediates a long-term response of the Arabidopsis thaliana circadian clock to sucrose. Proceedings of the National Academy of Sciences 108: 5104-5109

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

de Lucas M, Daviere JM, Rodriguez-Falcon M, Pontin M, Iglesias-Pedraz JM, Lorain S, Fankhauser C, Blazquez MA, Titarenko E, Prat S

(2008) A molecular framework for light and gibberellin control of cell elongation. Nature 451: 480-484

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Debast S, Nunes-Nesi A, Hajirezaei MR, Hofmann J, Sonnewald U, Fernie AR, Börnke F (2011) Altering trehalose-6-phosphate content in transgenic potato tubers affects tuber growth and alters responsiveness to hormones during sprouting. Plant Physiology 156: 1754-1771

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Delatte TL, Sedijani P, Kondou Y, Matsui M, de Jong GJ, Somsen GW, Wiese-Klinkenberg A, Primavesi LF, Paul MJ, Schluepmann H (2011) Growth arrest by trehalose-6-phosphate: An astonishing case of primary metabolite control over growth by way of the SnRK1 signaling pathway. Plant Physiology 157: 160

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Delorge I, Figueroa CM, Feil R, Lunn JE, van Dijck P (2015) Trehalose-6-phosphate synthase 1 is not the only active TPS in Arabidopsis thaliana. Biochemical Journal 466: 283-290

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ding Z, Doyle MR, Amasino RM, Davis SJ (2007) A complex genetic interaction between Arabidopsis thaliana TOC1 and CCA1/LHY in driving the circadian clock and in output regulation. Genetics 176: 1501-1510

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dobrenel T, Marchive C, Sormani R, Moreau M, Mozzo M, Montané M-H, Menand B, Robaglia C, Meyer C (2011) Regulation of plant growth and metabolism by the TOR kinase. Biochemical Society Transactions 39: 477

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dowson-Day MJ, Millar AJ (1999) Circadian dysfunction causes aberrant hypocotyl elongation patterns in Arabidopsis. The Plant Journal 17: 63-71

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Esmon CA, Tinsley AG, Ljung K, Sandberg G, Hearne LB, Liscum E (2006) A gradient of auxin and auxin-dependent transcription precedes tropic growth responses. Proceedings of the National Academy of Sciences of the United States of America 103: 236-241

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fu X, Richards DE, Ait-ali T, Hynes LW, Ougham H, Peng J, Harberd NP (2002) Gibberellin-mediated proteasome-dependent degradation of the barley DELLA protein SLN1 repressor. The Plant Cell 14: 3191-3200

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gangappa SN, Kumar SV (2017) DET1 and HY5 control PIF4-mediated thermosensory elongation growth through distinct mechanisms. Cell Reports 18: 344-351

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gendreau E, Traas J, Desnos T, Grandjean O, Caboche M, Hofte H (1997) Cellular basis of hypocotyl growth in Arabidopsis thaliana. Plant Physiology 114: 295-305

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Goda H, Sawa S, Asami T, Fujioka S, Shimada Y, Yoshida S (2004) Comprehensive comparison of auxin-regulated and brassinosteroid-regulated genes in Arabidopsis. Plant Physiology 134: 1555-1573

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gómez LD, Baud S, Gilday A, Li Y, Graham IA (2006) Delayed embryo development in the ARABIDOPSIS TREHALOSE-6-PHOSPHATE SYNTHASE 1 mutant is associated with altered cell wall structure, decreased cell division and starch accumulation. The Plant Journal

46: 69-84

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gómez LD, Gilday A, Feil R, Lunn JE, Graham IA (2010) AtTPS1-mediated trehalose 6-phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness to ABA in germinating seeds and stomatal guard cells. The Plant Journal 64: 1-13

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Graf A, Schlereth A, Stitt M, Smith AM (2010) Circadian control of carbohydrate availability for growth in Arabidopsis plants at night. Proceedings of the National Academy of Sciences 107: 9458-9463

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Griffiths CA, Sagar R, Geng Y, Primavesi LF, Patel MK, Passarelli MK, Gilmore IS, Steven RT, Bunch J, Paul MJ, Davis BG (2016) Chemical intervention in plant sugar signalling increases yield and resilience. Nature 540: 574-578

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Haydon MJ, Mielczarek O, Robertson FC, Hubbard KE, Webb AAR (2013) Photosynthetic entrainment of the Arabidopsis thaliana circadian clock. Nature 502: 689-692

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hayes S, Velanis CN, Jenkins GI, Franklin KA (2014) UV-B detected by the UVR8 photoreceptor antagonizes auxin signaling and plant shade avoidance. Proceedings of the National Academy of Sciences 111: 11894-11899

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jang JC, León P, Zhou L, Sheen J (1997) Hexokinase as a sugar sensor in higher plants. The Plant Cell 9: 5-19

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Keuskamp DH, Pollmann S, Voeselek LACJ, Peeters AJM, Pierik R (2010) Auxin transport through PIN-FORMED 3 (PIN3) controls shade avoidance and fitness during competition. Proceedings of the National Academy of Sciences 107: 22740-22744

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kim BC, Tennesen DJ, Last Robert L (1998) UV-B-induced photomorphogenesis in Arabidopsis thaliana. The Plant Journal 15: 667-674

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Koini MA, Avey L, Allen T, Tilley CA, Harberd NP, Whitelam GC, Franklin KA (2009) High temperature-mediated adaptations in plant architecture require the bHLH transcription factor PIF4. Current Biology 19: 408-413

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Koorneef M, Elgersma A, Hanhart CJ, van Loenen-Martinet EP, van Rijn L, Zeevaart JAD (1985) A gibberellin insensitive mutant of Arabidopsis thaliana. Physiologia Plantarum 65: 33-39

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kurata T, Yamamoto KT (1998) petit1, a conditional growth mutant of Arabidopsis defective in sucrose-dependent elongation growth. Plant Physiology 118: 793

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee DJ, Park JW, Lee HW, Kim J (2009) Genome-wide analysis of the auxin-responsive transcriptome downstream of iaa1 and its expression analysis reveal the diversity and complexity of auxin-regulated gene expression. Journal of Experimental Botany 60: 3935-3957

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Leivar P, Quail PH (2011) PIFs: pivotal components in a cellular signaling hub. Trends in Plant Science 16: 19-28

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Li Y, Darley CP, Ongaro V, Fleming A, Schipper O, Baldauf SL, McQueen-Mason SJ (2002) Plant expansins are a complex multigene family with an ancient evolutionary origin. Plant Physiology 128: 854-864

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Li Y, Van den Ende W, Rolland F (2014) Sucrose induction of anthocyanin biosynthesis is mediated by DELLA. Molecular Plant 7: 570-572

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lincoln C, Britton JH, Estelle M (1990) Growth and development of the axr1 mutants of Arabidopsis. Plant Cell 2: 1071-1080

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Liscum E, Hangarter RP (1991) Arabidopsis mutants lacking blue light-dependent inhibition of hypocotyl elongation. The Plant Cell 3: 685-694

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Liu Z, Zhang Y, Liu R, Hao H, Wang Z, Bi Y (2011) Phytochrome interacting factors (PIFs) are essential regulators for sucrose-induced hypocotyl elongation in Arabidopsis. Journal of Plant Physiology 168: 1771-1779

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lorrain S, Allen T, Duek PD, Whitelam GC, Fankhauser C (2008) Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. The Plant Journal 53: 312-323

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lunn JE, Feil R, Hendriks JHM, Gibon Y, Morcuende R, Osuna D, Sheible WR, Carillo P, Hajirezaei MR, Stitt M (2006) Sugar-induced increases in trehalose-6-phosphate are correlated with redox activation of ADPglucose pyrophosphorylase and higher rates of starch synthesis in Arabidopsis thaliana. Biochemical Journal 397: 139-148

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

MacGregor DR, Kendall SL, Florance H, Fedi F, Moore K, Paszkiewicz K, Smirnoff N, Penfield S (2015) Seed production temperature regulation of primary dormancy occurs through control of seed coat phenylpropanoid metabolism. New Phytologist 205: 642-652

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Mair A, Pedrotti L, Wurzinger B, Anrather D, Simeunovic A, Weiste C, Valerio C, Dietrich K, Kirchler T, Nägele T, Vicente Carbajosa J, Hanson J, Baena-González E, Chaban C, Weckwerth W, Dröge-Laser W, Teige M (2015) SnRK1-triggered switch of bZIP63 dimerization mediates the low-energy response in plants. eLife 4

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Más P, Alabadí D, Yanovsky MJ, Oyama T, Kay SA (2003) Dual role of TOC1 in the control of circadian and photomorphogenic responses in Arabidopsis. The Plant Cell 15: 223-236

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Matiolli CC, Tomaz JP, Duarte GT, Prado FM, Del Bem LEV, Silveira AB, Gauer L, Corrêa LGG, Drumond RD, Viana AJC, Di Mascio P, Meyer C, Vincentz M (2011) The Arabidopsis bZIP gene AtbZIP63 is a sensitive integrator of transient abscisic acid and glucose signals. Plant Physiology 157: 692-705

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Miyazaki Y, Jikumaru Y, Takase T, Saitoh A, Sugitani A, Kamiya Y, Kiyosue T (2016) Enhancement of hypocotyl elongation by LOV KELCH PROTEIN2 production is mediated by auxin and phytochrome-interacting factors in Arabidopsis thaliana. Plant Cell Reports 35:

455-467

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mizuno T, Nomoto Y, Oka H, Kitayama M, Takeuchi A, Tsubouchi M, Yamashino T (2014) Ambient temperature signal feeds into the circadian clock transcriptional circuitry through the EC night-time repressor in *Arabidopsis thaliana*. *Plant and Cell Physiology* 55: 958-976

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Moore B, Zhou L, Rolland F, Hall Q, Cheng W-H, Liu Y-X, Hwang I, Jones T, Sheen J (2003) Role of the *Arabidopsis* glucose sensor HXK1 in nutrient, light, and hormonal signaling. *Science* 300: 332-336

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nakamichi N, Kita M, Ito S, Yamashino T, Mizuno T (2005) PSEUDO-RESPONSE REGULATORS, PRR9, PRR7 and PRR5, together play essential roles close to the circadian clock of *Arabidopsis thaliana*. *Plant and Cell Physiology* 46: 686-698

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Noordally ZB, Ishii K, Atkins KA, Wetherill SJ, Kusakina J, Walton EJ, Kato M, Azuma M, Tanaka K, Hanaoka M, Dodd AN (2013) Circadian control of chloroplast transcription by a nuclear-encoded timing signal. *Science* 339: 1316-1319

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nozue K, Covington MF, Duek PD, Lorrain S, Fankhauser C, Harmer SL, Maloof JN (2007) Rhythmic growth explained by coincidence between internal and external cues. *Nature* 448: 358-361

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nuccio ML, Wu J, Mowers R, Zhou H-P, Meghji M, Primavesi LF, Paul MJ, Chen X, Gao Y, Haque E, Basu SS, Lagrimini LM (2015) Expression of trehalose-6-phosphate phosphatase in maize ears improves yield in well-watered and drought conditions. *Nat Biotech* 33: 862-869

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nunes C, O'Hara LE, Primavesi LF, Delatte TL, Schluepmann H, Somsen GW, Silva AB, Fevereiro PS, Wingler A, Paul MJ (2013) The trehalose 6-phosphate/SnRK1 signaling pathway primes growth recovery following relief of sink limitation. *Plant Physiology* 162: 1720-1732

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nunes C, Primavesi LF, Patel MK, Martinez-Barajas E, Powers SJ, Sagar R, Fevereiro PS, Davis BG, Paul MJ (2013) Inhibition of SnRK1 by metabolites: Tissue-dependent effects and cooperative inhibition by glucose 1-phosphate in combination with trehalose 6-phosphate. *Plant Physiology and Biochemistry* 63: 89-98

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nusinow DA, Helfer A, Hamilton EE, King JJ, Imaizumi T, Schultz TF, Farre EM, Kay SA (2011) The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* 475: 398-402

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Park S, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J, Rodrigues A, Chow TF, Alfred SE, Bonetta D, Finkelstein R, Provart NJ, Desveaux D, Rodriguez PL, McCourt P, Zhu JK, Schroeder JI, Volkman BF, Cutler SR (2009) Abscissic acid inhibits PP2Cs via the PYR/PYL family of ABA-binding START proteins. *Science* 324: 1068-1071

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Paul MJ, Jhurrea D, Zhang Y, Primavesi LF, Delatte T, Schluepmann H, Wingler A (2010) Up-regulation of biosynthetic processes associated with growth by trehalose 6-phosphate. *Plant Signaling & Behavior* 5: 386-392

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Penfield S, Rylott EL, Gilday AD, Graham S, Larson TR, Graham IA (2004) Reserve mobilization in the Arabidopsis endosperm fuels hypocotyl elongation in the dark, is independent of abscisic acid, and requires PHOSPHOENOLPYRUVATE CARBOXYKINASE1. The Plant Cell 16: 2705-2718

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Peng J, Carol P, Richards DE, King KE, Cowling RJ, Murphy GP, Harberd NP (1997) The Arabidopsis GAI gene defines a signaling pathway that negatively regulates gibberellin responses Genes & Development 11: 3194-3205

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ramon M, Rolland F, Thevelein JM, van Dijck P (2007) ABI4 mediates the effects of exogenous trehalose on Arabidopsis growth and starch breakdown. Plant Molecular Biology 63: 195-206

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Richardson K, Fowler S, Pullen C, Skelton C, Morris B, Putterill J (1998) T-DNA tagging of a flowering-time gene and improved gene transfer by in planta transformation of Arabidopsis. Functional Plant Biology 25: 125-130

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Salter MG, Franklin KA, Whitelam GC (2003) Gating of the rapid shade-avoidance response by the circadian clock in plants. Nature 426: 680-683

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schluepmann H, Pellny T, van Dijken A, Smekens S, Paul M (2003) Trehalose 6-phosphate is indispensable for carbohydrate utilization and growth in Arabidopsis thaliana. Proceedings of the National Academy of Sciences of the United States of America 100: 6849-6854

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shin J, Sánchez-Villarreal A, Davis AM, Du S-x, Berendzen KW, Koncz C, Ding Z, Li C, Davis SJ (2017) The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner. Plant, Cell & Environment 40: 997-1008

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Smekens S, Ma J, Hanson J, Rolland F (2010) Sugar signals and molecular networks controlling plant growth. Current Opinion in Plant Biology 13: 273-278

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stewart JL, Maloof JN, Nemhauser JL (2011) PIF genes mediate the effect of sucrose on seedling growth dynamics. PLoS One 6: e19894

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stewart Lilley JL, Gee CW, Sairanen I, Ljung K, Nemhauser JL (2012) An endogenous carbon-sensing pathway triggers increased auxin flux and hypocotyl elongation. Plant Physiology 160: 2261-2270

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sulpice R, Flis A, Ivakov AA, Apelt F, Krohn N, Encke B, Abel C, Feil R, Lunn JE, Stitt M (2014) Arabidopsis coordinates the diurnal regulation of carbon allocation and growth across a wide range of photoperiods. Molecular Plant 7: 137-155

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takahashi F, Sato-Nara K, Kobayashi K, Suzuki M, Suzuki H (2003) Sugar-induced adventitious roots in Arabidopsis seedlings. Journal of Plant Research 116: 83-91

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vandesteene L, Ramon M, Le Roy K, Van Dijck P, Rolland F (2010) A single active trehalose-6-P synthase (TPS) and a family of putative

regulatory TPS-like proteins in Arabidopsis. Molecular Plant 3: 406-419

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M (2013) Regulation of flowering by trehalose-6-phosphate signaling in Arabidopsis thaliana. Science 339: 704-707

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang Z-Y, Tobin EM (1998) Constitutive expression of the CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own expression. Cell 93: 1207-1217

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Weinig C, Johnston JA, Willis CG, Maloof JN (2007) Antagonistic multilevel selection on size and architecture in variable density settings. Evolution 61: 58-67

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wigge PA (2013) Ambient temperature signalling in plants. Current Opinion in Plant Biology 16: 661-666

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wingler A, Delatte TL, O'Hara LE, Primavesi LF, Jhurreea D, Paul MJ, Schluepmann H (2012) Trehalose 6-phosphate is required for the onset of leaf senescence associated with high carbon availability. Plant Physiology 158: 1241-1251

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ (2007) An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets. PLoS One 2: e718

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yadav UP, Ivakov A, Feil R, Duan GY, Walther D, Gialalisco P, Piques M, Carillo P, Hubberten H-M, Stitt M, Lunn JE (2014) The sucrose-trehalose 6-phosphate (Tre6P) nexus: specificity and mechanisms of sucrose signalling by Tre6P. Journal of Experimental Botany 65: 1051-1068

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yamamoto Y, Sato E, Shimizu T, Nakamichi N, Sato S, Kato T, Tabata S, Nagatani A, Yamashino T, Mizuno T (2003) Comparative genetic studies on the APRR5 and APRR7 genes belonging to the APRR1/TOC1 quintet implicated in circadian rhythm, control of flowering time, and early photomorphogenesis. Plant and Cell Physiology 44: 1119-1130

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang Y, Liu Z, Wang J, Chen Y, Bi Y, He J (2015) Brassinosteroid is required for sugar promotion of hypocotyl elongation in Arabidopsis in darkness. Planta 242: 881-893

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang Y, Liu Z, Wang L, Zheng S, Xie J, Bi Y (2010) Sucrose-induced hypocotyl elongation of Arabidopsis seedlings in darkness depends on the presence of gibberellins. Journal of Plant Physiology 167: 1130-1136

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang Y, Primavesi LF, Jhurreea D, Andralojc PJ, Mitchell RAC, Powers SJ, Schluepmann H, Delatte T, Wingler A, Paul MJ (2009) Inhibition of SNF1-related protein kinase 1 activity and regulation of metabolic pathways by trehalose-6-phosphate. Plant Physiology 149: 1860-1871

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang Y, Shewry PR, Jones H, Barcelo P, Lazzeri PA, Halford NG (2001) Expression of antisense SnRK1 protein kinase sequence causes abnormal pollen development and male sterility in transgenic barley. The Plant Journal 28: 431-441

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang Z, Zhu J-Y, Roh J, Marchive C, Kim S-K, Meyer C, Sun Y, Wang W, Wang Z-Y (2016) TOR signaling promotes accumulation of BZR1 to balance growth with carbon availability in Arabidopsis. Current Biology 26: 1854-1860

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)